

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C12N 15/12, 9/64, A61K 38/48, G01N 33/50</b>		<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 98/31801</b> <b>(43) International Publication Date:</b> 23 July 1998 (23.07.98)
<b>(21) International Application Number:</b> PCT/US98/00969 <b>(22) International Filing Date:</b> 21 January 1998 (21.01.98)  <b>(30) Priority Data:</b> 60/034,205                      21 January 1997 (21.01.97)                      US 60/054,800                      5 August 1997 (05.08.97)                      US  <b>(71) Applicants:</b> HUMAN GENOME SCIENCES, INC. [US/US]; 9410 Key West Avenue, Rockville, MD 20850 (US). THE REGENTS OF THE UNIVERSITY OF MICHIGAN [US/US]; Wolverine Tower, Room 2071, 3003 South State Street, Ann Arbor, MI 48109-1280 (US).  <b>(72) Inventors:</b> NI, Jian; 5502 Manorfield Road, Rockville, MD 20853 (US). ROSEN, Craig, A.; 22400 Rolling Hill Road, Laytonsville, MD 20882 (US). DIXIT, Vishva, M.; 26750 Shady Oaks Court, Los Altos Hills, CA 94022 (US). GENTZ, Reiner, L.; 13404 Fairland Park Drive, Silver Spring, MD 20904 (US). KENNY, Joseph, J.; Apartment H12, 2014 Baltimore Road, Rockville, MD 20851 (US).  <b>(74) Agents:</b> STEFFE, Eric, K.; Sterne, Kessler, Goldstein & Fox P.L.L.C., Suite 600, 1100 New York Avenue, N.W., Washington, DC 20005-3934 (US) et al.			<b>(81) Designated States:</b> CA, JP, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i> <i>With an indication in relation to a deposited microorganism furnished under Rule 13bis separately from the description.</i> <i>Date of receipt by the International Bureau:</i> 20 May 1998 (20.05.1998)
<b>(54) Title:</b> I-FLICE, A NOVEL INHIBITOR OF TUMOR NECROSIS FACTOR RECEPTOR-1 AND CD-95 INDUCED APOPTOSIS			
<b>(57) Abstract</b>  The present invention relates to a novel I-FLICE-1 or I-FLICE-2 protein which is a novel inhibitor of TNFR-1 and CD-95 induced apoptosis. In particular, isolated nucleic acid molecules are provided encoding the human I-FLICE-1 or I-FLICE-2 protein. I-FLICE-1 or I-FLICE-2 polypeptides are also provided as are vectors, host cells and recombinant methods for producing the same. The invention further relates to screening methods for identifying agonists and antagonists of I-FLICE-1 or I-FLICE-2 activity. Also provided are therapeutic methods for treating diseases and disorders associated with apoptosis.			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## I-FLICE, A NOVEL INHIBITOR OF TUMOR NECROSIS FACTOR RECEPTOR-1 AND CD-95 INDUCED APOPTOSIS

### *Background of the Invention*

#### *Field of the Invention*

5           The present invention relates to a novel inhibitor of TNFR-1 and CD-95 induced apoptosis. More specifically, isolated nucleic acid molecules are provided encoding a human I-FLICE (Inhibitor of FLICE (FADD-like ICE)) polynucleotides. I-FLICE polypeptides are also provided, as are vectors, host cells and recombinant methods for producing the same. The invention further  
10 relates to screening methods for identifying agonists and antagonists of I-FLICE activity. Also provided are therapeutic methods for treating diseases and disorders associated with apoptosis.

#### *Related Art*

15           The cell death machinery is conserved throughout evolution and is composed of activators, inhibitors, and effectors (Chinnaiyan, A.M. and Dixit, V.M., *Curr. Biol.* 6:555-562 (1996)). The effector arm of the cell death pathway is composed of a rapidly growing family of cysteine aspartate-specific proteases termed caspases (Alnemri, E.S., et al., *Cell* 87:171 (1996)). As implied by the  
20 name, these cysteine proteases cleave substrates following an aspartate residue (Alnemri, E.S., et al., *Cell* 87:171 (1996); Walker, N.P., et al., *Cell* 78:343-352 (1994)). Caspases are normally present as single polypeptide zymogens and contain an amino-terminal prodomain, and large and small catalytic subunits (Wilson, K.P., et al., *Nature* 370:270-274 (1994); Rotonda, J., et al., *Nat. Struct.*  
25 *Biol.* 3:619-625 (1996); Fraser, A. and Evan, G., *Cell* 85:781-784 (1996)). The two chain active enzyme (composed of the large and small subunits) is obtained following proteolytic processing at internal Asp residues (Wilson, K.P., et al., *Nature* 370:270-274 (1994); Rotonda, J., et al., *Nat. Struct. Biol.* 3:619-625 (1996); Fraser, A. and Evan, G., *Cell* 85:781-784 (1996)). As such, caspases are  
30 capable of activating each other in a manner analogous to zymogen activation that

-2-

is observed in the coagulation cascade (Boldin, M.P., et al., *Cell* 85:805-815 (1996)). The identification of FLICE and Mch4/FLICE2 as receptor associated caspases suggested a surprisingly direct mechanism for activation of the death pathway by the cytotoxic receptors CD-95 and TNFR-1 (Boldin, M.P., et al., *Cell* 85:805-815 (1996); Muzio, M., et al., *Cell* 85:817-827 (1996); Vincenz, C. and Dixit, V.M., *J. Biol. Chem.* 272:6578-6583 (1997); Chinnaiyan, A.M., et al., *Cell* 81:505-512 (1995)). Upon activation, both receptors use their death domains to bind the corresponding domain in the adaptor molecule FADD (Fas-associated death domain protein) (Muzio, M., et al., *Cell* 85:817-827 (1996); Vincenz, C. and Dixit, V.M., *J. Biol. Chem.* 272:6578-6583 (1997); Chinnaiyan, A.M., et al., *Cell* 81:505-512 (1995)). Dominant negative versions of FADD that lack the N-terminal segment but still retain the death domain potently inhibit both CD-95 and TNFR-1 induced apoptosis (Chinnaiyan, A.M., et al., *J. Biol. Chem.* 271:4961-4965 (1996); Muzio, M., et al., *J. Biol. Chem.* 272:2952-2956 (1997)). Given the importance of the N-terminal segment in engaging the death pathway, it has been termed the death effector domain (DED) (Chinnaiyan, A.M., et al., *J. Biol. Chem.* 271:4961-4965 (1996)).

Remarkably, the DED is present within the prodomain of FLICE and Mch4/FLICE2 and mutagenesis studies suggest that a homophilic interaction between the DED of FADD and the corresponding domain in FLICE or Mch4/FLICE2 is responsible for the recruitment of these proteases to the CD-95 and TNFR-1 signalling complexes (Muzio, M., et al., *Cell* 85:817-827 (1996); Vincenz, C. and Dixit, V.M., *J. Biol. Chem.* 272:6578-6583 (1997); Chinnaiyan, A.M., et al., *Cell* 81:505-512 (1995); Chinnaiyan, A.M., et al., *J. Biol. Chem.* 271:4961-4965 (1996)). Taken together, these data are consistent with FLICE and Mch4/FLICE2 being apical enzymes that initiate precipitous proteolytic processing of downstream caspases resulting in apoptosis (Boldin, M.P., et al., *Cell* 85:805-815 (1996); Srinivasula, S.M., et al., *PNAS* 93:14486-14491 (1996); Fernandes-Alnemri, T., et al., *PNAS* 93:7464-7469 (1996); Henkart, P.A., *Immunity* 4:195-201 (1996)). A number of viral gene products antagonize CD-95



and TNFR-1 mediated killing as a means to persist in the infected host (Shen, Y. and Shenk, T.S., *Current Opinion in Genetics and Development* 5:105-111 (1995)). The poxvirus encoded serpin CrmA and baculovirus gene product p35 are direct caspase inhibitors (Walker, N.P., et al., *Cell* 78:343-352 (1994)). In contrast, the molluscum contagiosum virus protein MC159 and the equine herpes virus protein E8 encode DED-containing decoy molecules that bind to either FADD (MC159) or FLICE (E8) and disrupt assembly of the receptor signalling complex, thereby abrogating the death signal (Hu, S., et al., *J. Biol. Chem.* 272:9621-9624 (1997); Bertin, J., et al., *PNAS* 94:1172-1176 (1997); Thome, M., et al., *Nature* 386:527-521 (1997)). The existence of these viral inhibitors has raised the question of whether functionally equivalent molecules are encoded in the mammalian genome.

There is a need for factors, such as the polypeptides of the present invention, that are useful for inhibiting apoptosis for therapeutic purposes, for example, in the treatment of Alzheimer's disease, Parkinson's disease, rheumatoid arthritis, septic shock, sepsis, stroke, CNS inflammation, osteoporosis, ischemia, reperfusion injury, cell death associated with cardiovascular disease, polycystic kidney disease, apoptosis of endothelial cells in cardiovascular disease, degenerative liver disease, MS and head injury damage. There is a need, therefore, for the identification and characterization of such factors that are inhibitors of apoptosis, such as the I-FLICE-1 and I-FLICE-2 polypeptides of the present invention, which can play a role in preventing, ameliorating or correcting the diseases and disorders associated with apoptosis.

### ***Summary of the Invention***

The present invention provides isolated nucleic acid molecules comprising a polynucleotide encoding the I-FLICE-1 polypeptide having the amino acid sequence shown in SEQ ID NO:2 or the amino acid sequence encoded by the cDNA clone deposited in a bacterial host as ATCC Deposit Number 209038 on

May 15, 1997. The present invention provides isolated nucleic acid molecules comprising a polynucleotide encoding the I-FLICE-2 polypeptide having the amino acid sequence shown in SEQ ID NO:6 or the amino acid sequence encoded by the cDNA clone deposited in a bacterial host as ATCC Deposit Number 209041 on May 15, 1997.

The present invention also relates to recombinant vectors, which include the isolated nucleic acid molecules of the present invention, and to host cells containing the recombinant vectors, as well as to methods of making such vectors and host cells and for using them for production of I-FLICE-1 or I-FLICE-2 polypeptides or peptides by recombinant techniques.

The invention further provides an isolated I-FLICE-1 or I-FLICE-2 polypeptides having an amino acid sequence encoded by the polynucleotides described herein.

The invention further provides a diagnostic method useful during diagnosis or prognosis of a disease states resulting from aberrant cell proliferation due to alterations in I-FLICE-1 or I-FLICE-2 expression.

The present invention also provides a screening method for determining whether a candidate agonist or antagonist is capable of enhancing or inhibiting a cellular activity of either an I-FLICE-1 or I-FLICE-2 polypeptide. The method involves contacting cells which express one or both of the I-FLICE-1 or I-FLICE-2 polypeptides with a candidate compound, assaying a cellular response, and comparing the cellular response to a standard cellular response, the standard being assayed in absence of the candidate compound, whereby an increased cellular response over the standard indicates that the candidate compound is an agonist of the polypeptide activity and a decreased cellular response compared to the standard indicates that the candidate compound is an antagonist of the activity.

An additional aspect of the invention is related to a method for treating an individual in need of an increased level of I-FLICE-1 or I-FLICE-2 activity in the body comprising administering to such an individual a composition comprising

a therapeutically effective amount of an isolated I-FLICE-1 or I-FLICE-2 polypeptide of the invention or an agonist thereof.

A still further aspect of the invention is related to a method for treating an individual in need of a decreased level of I-FLICE-1 or I-FLICE-2 activity in the body comprising, administering to such an individual a composition comprising a therapeutically effective amount of an I-FLICE-1 or I-FLICE-2 antagonist.

### *Brief Description of the Figures*

FIG. 1A-1B shows the nucleotide (SEQ ID NO:1) and deduced amino acid (SEQ ID NO:2) sequences of I-FLICE-1 (HSLAZ11). The protein has 480 amino acid residues and a deduced molecular weight of about 55.3 kDa.

FIG. 2 shows the regions of similarity between the amino acid sequences of the I-FLICE-1, I-FLICE-2, FLICE (SEQ ID NO:3), and Mch4 (SEQ ID NO:4). Shading (with solid black) indicates residues that match the consensus sequence exactly.

FIG. 3 shows an analysis of the I-FLICE-1 amino acid sequence. Alpha, beta, turn and coil regions; hydrophilicity and hydrophobicity; amphipathic regions; flexible regions; antigenic index and surface probability are shown. In the "Antigenic Index-Jameson-Wolf" graph, amino acid residues about 41 to about 92, about 155 to about 249, about 332 to about 447 in FIG. 1A-1B (SEQ ID NO:2) correspond to the shown highly antigenic regions of the I-FLICE-1 protein.

FIG. 4A-4C shows the nucleotide (SEQ ID NO:5) and deduced amino acid (SEQ ID NO:6) sequences of I-FLICE-2 (HCEBJ50). The protein has 348 amino acid residues and a deduced molecular weight of about 39.2 kDa.

FIG. 5 shows an analysis of the I-FLICE-2 amino acid sequence. Alpha, beta, turn and coil regions; hydrophilicity and hydrophobicity; amphipathic regions; flexible regions; antigenic index and surface probability are shown. In the "Antigenic Index - Jameson-Wolf" graph, amino acid residues about 62 to

about 136, about 184 to about 193, about 205 to about 341 in FIG. 4A-4C (SEQ ID NO:6) correspond to the shown highly antigenic regions of the I-FLICE-2 protein.

FIG. 6A-6B shows I-FLICE-1 inhibition of apoptosis. Overexpression of I-FLICE-1 attenuated TNFR-1 (panel A) and CD-95 (panel B) induced cell death. 293 (panel A) or 293-EBNA (panel B) cells were co-transfected with the indicated plasmids together with the reporter construct pCMV  $\beta$ -galactosidase. The data shown are the percentage of blebbing blue cells as a function of total number of blue cells counted.

### *Detailed Description*

The present invention provides isolated nucleic acid molecules comprising a polynucleotide encoding an I-FLICE-1 or I-FLICE-2 polypeptide having the amino acid sequence shown in SEQ ID NO:2 or SEQ ID NO:6, respectively, which was determined by sequencing a cloned cDNA. The I-FLICE-1 protein of the present invention shares sequence homology with FLICE and Mch4 (FIG. 2) (SEQ ID NOs:3 and 4). The nucleotide sequence shown in SEQ ID NO:1 was obtained by sequencing a cDNA clone (HSLAZ11), which was deposited on May 15, 1997 at the American Type Culture Collection, 12301 Park Lawn Drive, Rockville, Maryland 20852, and given accession number 209038. The deposited clone is inserted in the pBluescript SK(-) plasmid (Stratagene, La Jolla, CA) using the EcoRI and XhoI restriction endonuclease cleavage sites. The I-FLICE-2 protein of the present invention shares sequence homology with FLICE and Mch4 (FIG. 2 (SEQ ID NOs:3 and 4)). The nucleotide sequence shown in SEQ ID NO:5 was obtained by sequencing a cDNA clone (HCEBJ50), which was deposited on May 15, 1997 at the American Type Culture Collection, 12301 Park Lawn Drive, Rockville, Maryland 20852, and given accession number 209041. The deposited clone is inserted in the pBluescript SK(-) plasmid (Stratagene, La Jolla, CA) using the EcoRI and XhoI restriction endonuclease cleavage sites.

### *Nucleic Acid Molecules*

Unless otherwise indicated, all nucleotide sequences determined by sequencing a DNA molecule herein were determined using an automated DNA sequencer (such as the Model 373 from Applied Biosystems, Inc.), and all amino acid sequences of polypeptides encoded by DNA molecules determined herein were predicted by translation of a DNA sequence determined as above. Therefore, as is known in the art for any DNA sequence determined by this automated approach, any nucleotide sequence determined herein may contain some errors. Nucleotide sequences determined by automation are typically at least about 90% identical, more typically at least about 95% to at least about 99.9% identical to the actual nucleotide sequence of the sequenced DNA molecule. The actual sequence can be more precisely determined by other approaches including manual DNA sequencing methods well known in the art. As is also known in the art, a single insertion or deletion in a determined nucleotide sequence compared to the actual sequence will cause a frame shift in translation of the nucleotide sequence such that the predicted amino acid sequence encoded by a determined nucleotide sequence will be completely different from the amino acid sequence actually encoded by the sequenced DNA molecule, beginning at the point of such an insertion or deletion.

Using the information provided herein, such as the nucleotide sequence in SEQ ID NO:1 or SEQ ID NO:5, a nucleic acid molecule of the present invention encoding an I-FLICE-1 or I-FLICE-2 polypeptide may be obtained using standard cloning and screening procedures, such as those for cloning cDNAs using mRNA as starting material. Illustrative of the invention, the nucleic acid molecule described in SEQ ID NO:1 was discovered in a cDNA library derived from human umbilical vein endothelial cell. The gene was also identified in cDNA libraries from smooth muscle. The determined nucleotide sequence of the I-FLICE-1 cDNA of SEQ ID NO:1 contains an open reading frame encoding a protein of about 480 amino acid residues and a deduced

molecular weight of about 55.3 kDa. The I-FLICE-1 protein shown in SEQ ID NO:2 is overall about 29% identical and about 54% similar to FLICE (FIG. 2 (SEQ ID NO:3)).

Also illustrative of the invention, the nucleic acid molecule described in  
5 SEQ ID NO:5 was discovered in a cDNA library derived from human umbilical vein endothelial cell. The gene was also identified in cDNA libraries from brain tissue isolated from the cerebellum. The determined nucleotide sequence of the I-FLICE-2 cDNA of SEQ ID NO:5 contains an open reading frame encoding a protein of about 348 amino acid residues and a deduced molecular weight of  
10 about 39 kDa. The I-FLICE-2 protein shown in SEQ ID NO:6 is overall about 28% identical and about 54% similar to FLICE (FIG. 2 (SEQ ID NO:3)).

In addition, I-FLICE-1 and I-FLICE-2 are nearly identical over the majority of their sequences; however, I-FLICE-1 has additional amino acids comprising the N-terminal region of the protein. The amino terminal domains of  
15 both I-FLICE-1 and I-FLICE-2 exhibit significant sequence similarity to the DED domain of the FADD protein (Hu, S. et al., *J. Biol. Chem.* 272:17255-17257 (1997); Irmeler, M., et al., *Nature* 388:190-195 (1997)), the domain through which FLICE proteins and death receptors interact. The amino terminal domain of I-FLICE-2 consists of only a single DED/FADD homology domain (comprising  
20 amino acid residues from about 1 to about 75 in SEQ ID NO:6), while the additional amino acids found in the amino terminal domain of I-FLICE-1 appear to provide a second DED/FADD homology domain (comprising amino acid residues from about 1 to about 75 and amino acids residues from about 91 to about 171 in SEQ ID NO:2). The carboxy terminal domains of the both I-FLICE-  
25 1 and I-FLICE-2 also contain significant sequence similarity to the active subunit domains of the ICE/CED-3 family of cysteine proteases (amino acids residues from about 172 to about 375 and amino acid residues from about 376 to about 480 in SEQ ID NO:2; amino acids residues from about 76 to about 252 and amino acid residues from about 253 to about 348 in SEQ ID NO:6). Neither I-  
30 FLICE-1 or I-FLICE-2 contain the catalytic cysteine that is normally embedded

in the conserved pentapeptide QACRG or QACQG motif present in all known caspases. Rather, both I-FLICE-1 and I-FLICE-2 have the pentapeptide sequence QNYVV (amino acid residues from about 358 to about 362 in SEQ ID NO:2 and amino acid residues from about 244 to about 248 in SEQ ID NO:6). Further, only  
5 three of seven conserved residues that form the substrate binding pocket found in all caspases are present in I-FLICE-1 and I-FLICE-2. Given the lack of conservation of key residues involved in catalysis and substrate binding, it can be concluded that I-FLICE-1 and I-FLICE-2 are not cysteine proteases and are incapable of substrate binding, thus, providing these proteins with a dominant  
10 negative inhibitory function. I-FLICE-1 and I-FLICE-2 are the first examples of catalytically inert caspases that can inhibit apoptosis.

As indicated, nucleic acid molecules of the present invention may be in the form of RNA, such as mRNA, or in the form of DNA, including, for instance, cDNA and genomic DNA obtained by cloning or produced synthetically. The  
15 DNA may be double-stranded or single-stranded. Single-stranded DNA or RNA may be the coding strand, also known as the sense strand, or it may be the non-coding strand, also referred to as the anti-sense strand.

By "isolated" nucleic acid molecule(s) is intended a nucleic acid molecule, DNA or RNA, which has been removed from its native environment. For  
20 example, recombinant DNA molecules contained in a vector are considered isolated for the purposes of the present invention. Further examples of isolated DNA molecules include recombinant DNA molecules maintained in heterologous host cells or purified (partially or substantially) DNA molecules in solution. Isolated RNA molecules include *in vivo* or *in vitro* RNA transcripts of the DNA  
25 molecules of the present invention. Isolated nucleic acid molecules according to the present invention further include such molecules produced synthetically.

Isolated nucleic acid molecules of the present invention include DNA molecules comprising an open reading frame (ORF) shown in SEQ ID NO:1 or  
30 EQ ID NO:5; DNA molecules comprising the coding sequence for the I-FLICE -1 or I-FLICE-2 protein; and DNA molecules which comprise a sequence

substantially different from those described above but which, due to the degeneracy of the genetic code, still encode the I-FLICE-1 or I-FLICE-2 protein. Of course, the genetic code is well known in the art. Thus, it would be routine for one skilled in the art to generate such degenerate variants.

5 In addition, the invention provides nucleic acid molecules having nucleotide sequences related to extensive portions of SEQ ID NO:1 which have been determined from the following related cDNA clones: HOSBY07R (SEQ ID NO:23), HSAVA13R (SEQ ID NO:24), HLFBD88R (SEQ ID NO:25), HOSAH65R (SEQ ID NO:26), HUVBS23R (SEQ ID NO:27), HHFFJ01RA (SEQ ID NO:28), HUVBL22R (SEQ ID NO:29), and HUVBX15R (SEQ ID NO:30).

10 The invention also provides nucleic acid molecules having nucleotide sequences related to extensive portions of SEQ ID NO:5 (I-FLICE-2) which have been determined from the following related cDNA clones: HTNBE58R (SEQ ID NO:31), HTPBE58R (SEQ ID NO:32), HOSBY07R (SEQ ID NO:23), HSAVA13R (SEQ ID NO:24), HLFBD88R (SEQ ID NO:25), HOSAH65R (SEQ ID NO:26), and HHFFJ01RA (SEQ ID NO:28).

15 In another aspect, the invention provides isolated nucleic acid molecules encoding the I-FLICE-1 polypeptide having an amino acid sequence as encoded by the cDNA clone contained in the plasmid deposited as ATCC Deposit No. 209038 on May 15, 1997. The invention also provides isolated nucleic acid molecules encoding the I-FLICE-2 polypeptide having an amino acid sequence as encoded by the cDNA clone contained in the plasmid deposited as ATCC Deposit No. 209041 on May 15, 1997. In a further embodiment, nucleic acid molecules are provided encoding the I-FLICE-1 or I-FLICE-2 polypeptide or the full-length I-FLICE polypeptide lacking the N-terminal methionine. The invention also provides an isolated nucleic acid molecule having the nucleotide sequence shown in SEQ ID NO:1 or SEQ ID NO:5 or the nucleotide sequence of the I-FLICE-1 or I-FLICE-2 cDNA contained in the above-described deposited clones, or a nucleic acid molecule having a sequence complementary to one of the

20

25

30



-11-

above sequences. Such isolated molecules, particularly DNA molecules, are useful as probes for gene mapping, by *in situ* hybridization with chromosomes, and for detecting expression of the I-FLICE-1 or I-FLICE-2 gene in human tissue, for instance, by Northern blot analysis.

5           The present invention is further directed to fragments of the isolated nucleic acid molecules described herein. By a fragment of an isolated nucleic acid molecule having the nucleotide sequence of the deposited cDNA or the nucleotide sequence shown in SEQ ID NO:1 or SEQ ID NO:5 is intended  
10 fragments at least about 15 nt, and more preferably at least about 20 nt, still more preferably at least about 30 nt, and even more preferably, at least about 40 nt in length which are useful as diagnostic probes and primers as discussed herein. Of course larger DNA fragments 50, 100, 150, 200, 250, 300, 350, 400, 450, 500, 550, 600, 650, 700, 750, 800, 850, 900, 950, 1000, 1050, 1100, 1150, 1200, 1250, 1300, 1350, 1400, 1450, 1500, 1550, 1600, 1650, 1700, 1750, 1800, 1850, 1900,  
15 1950, 2000, or 2016 nt in length of the sequence shown in SEQ ID NO:1 are also useful according to the present invention as are fragments corresponding to most, if not all, of the nucleotide sequence of the cDNA clone contained in the plasmid deposited as ATCC Deposit No. 209038 or as shown in SEQ ID NO:1. Similarly, larger DNA fragments 50, 100, 150, 200, 250, 300, 350, 400, 450, 500, 550, 600,  
20 650, 700, 750, 800, 850, 900, 950, 1000, 1050, 1100, 1150, 1200, 1250, 1300, 1350, 1400, 1450, 1500, 1550, 1600, 1650, 1700, 1750, 1800, 1850, 1900, 1950, 2000, 2050, 2100, 2150, 2200, 2250, 2300, 2350, 2400, 2450, 2500, or 2547 nt in length of the sequence shown in SEQ ID NO:5 are also useful according to the present invention as are fragments corresponding to most, if not all, of the  
25 nucleotide sequence of the cDNA clone contained in the plasmid deposited as ATCC Deposit No. 209041 or as shown in SEQ ID NO:5. By a fragment at least 20 nt in length, for example, is intended fragments which include 20 or more contiguous bases from the nucleotide sequence of the deposited cDNA or the nucleotide sequence as shown in SEQ ID NO:1 or SEQ ID NO:5.

In a more specific embodiment, the nucleic acid molecules of the present invention do not include the sequences, nucleic acid molecules (*e.g.*, clones), or nucleic acid inserts identified in one or more of the following GenBank Accession Reports: AA001257, AA151642, AA149562, C05730, AA565691, AA467756, D83882, AA002262, AA115793, AA467995, AA115792, AA467938, W60406, AA358042, AA468056, W23795, AA358043, T93307, AA453850, AA379905, AA296229, H15978, AA501289, AA296309, AA296174, T30922, T48754, AA453766, C05795, AA198928, N94588, H15052, Z42895, F13176, W52946, AA558404, AA070614, AA613966, AA525331, AA663074, AA135811, AA526099, AA302978, H68343, AA610255, AA229005, T05118, T30864, AA302968, or AA364006, all of which are incorporated herein by reference.

Preferred nucleic acid fragments of the present invention include nucleic acid molecules encoding epitope-bearing portions of the I-FLICE-1 protein. In particular, such nucleic acid fragments of the present invention include nucleic acid molecules encoding: a polypeptide comprising amino acid residues from about 41 to about 92 in SEQ ID NO:2; a polypeptide comprising amino acid residues from about 155 to about 249 in SEQ ID NO:2; a polypeptide comprising amino acid residues from about 332 to about 474 in SEQ ID NO:2. The inventors have determined that the above polypeptide fragments are antigenic regions of the I-FLICE-1 protein. Methods for determining other such epitope-bearing portions of the I-FLICE-1 protein are described in detail below.

Preferred nucleic acid fragments of the present invention include nucleic acid molecules encoding epitope-bearing portions of the I-FLICE-2 protein. In particular, such nucleic acid fragments of the present invention include nucleic acid molecules encoding: a polypeptide comprising amino acid residues from about 62 to about 136 in SEQ ID NO:6; a polypeptide comprising amino acid residues from about 184 to about 193 in SEQ ID NO:6; a polypeptide comprising amino acid residues from about 205 to about 341 in SEQ ID NO:6. The inventors have determined that the above polypeptide fragments are antigenic regions of the

I-FLICE-2 protein. Methods for determining other such epitope-bearing portions of the I-FLICE-2 protein are described in detail below.

In another aspect, the invention provides an isolated nucleic acid molecule comprising a polynucleotide which hybridizes under stringent hybridization conditions to a portion of the polynucleotide in a nucleic acid molecule of the invention described above, for instance, the cDNA clones contained in ATCC Deposit 209038 or ATCC Deposit 209041. By "stringent hybridization conditions" is intended overnight incubation at 42°C in a solution comprising: 50% formamide, 5x SSC (150 mM NaCl, 15mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 g/ml denatured, sheared salmon sperm DNA, followed by washing the filters in 0.1x SSC at about 65°C.

By a polynucleotide which hybridizes to a "portion" of a polynucleotide is intended a polynucleotide (either DNA or RNA) hybridizing to at least about 15 nucleotides (nt), and more preferably at least about 20 nt, still more preferably at least about 30 nt, and even more preferably about 30-70 nt of the reference polynucleotide. These are useful as diagnostic probes and primers as discussed above and in more detail below.

By a portion of a polynucleotide of "at least 20 nt in length," for example, is intended 20 or more contiguous nucleotides from the nucleotide sequence of the reference polynucleotide (e.g., the deposited cDNAs or the nucleotide sequence as shown in SEQ ID NO:1 or SEQ ID NO:5). Of course, a polynucleotide which hybridizes only to a poly A sequence (such as the 3' terminal poly(A) tract of the I-FLICE-1 cDNA shown in SEQ ID NO:1 or the I-FLICE-2 cDNA shown in SEQ ID NO:5), or to a complementary stretch of T (or U) residues, would not be included in a polynucleotide of the invention used to hybridize to a portion of a nucleic acid of the invention, since such a polynucleotide would hybridize to any nucleic acid molecule containing a poly (A) stretch or the complement thereof (e.g., practically any double-stranded cDNA clone).

As indicated, nucleic acid molecules of the present invention which encode an I-FLICE-1 or I-FLICE-2 polypeptide may include, but are not limited to those encoding the amino acid sequence of the polypeptide, by itself; the coding sequence for the mature polypeptide and additional sequences, such as those encoding a secretory sequence, such as a pre-, or pro- or prepro- protein sequence; the coding sequence of the polypeptide, with or without the aforementioned additional coding sequences, together with additional, non-coding sequences, including for example, but not limited to introns and non-coding 5' and 3' sequences, such as the transcribed, non-translated sequences that play a role in transcription, mRNA processing, including splicing and polyadenylation signals, for example - ribosome binding and stability of mRNA; an additional coding sequence which codes for additional amino acids, such as those which provide additional functionalities. Thus, the sequence encoding the polypeptide may be fused to a marker sequence, such as a sequence encoding a peptide which facilitates purification of the fused polypeptide. In certain preferred embodiments of this aspect of the invention, the marker amino acid sequence is a hexa-histidine peptide, such as the tag provided in a pQE vector (Qiagen, Inc.), among others, many of which are commercially available. As described in Gentz *et al.*, *Proc. Natl. Acad. Sci. USA* 86:821-824 (1989), for instance, hexa-histidine provides for convenient purification of the fusion protein. The "HA" tag is another peptide useful for purification which corresponds to an epitope derived from the influenza hemagglutinin protein, which has been described by Wilson *et al.*, *Cell* 37:767-778 (1984). As discussed below, other such fusion proteins include the I-FLICE-1 or I-FLICE-2 fused to Fc at the N- or C-terminus.

The present invention further relates to variants of the nucleic acid molecules of the present invention, which encode portions, analogs or derivatives of the I-FLICE-1 or I-FLICE-2 protein. Variants may occur naturally, such as a natural allelic variant. By an "allelic variant" is intended one of several alternate forms of a gene occupying a given locus on a chromosome of an organism.

*Genes II*, Lewin, B., ed., John Wiley & Sons, New York (1985). Non-naturally occurring variants may be produced using art-known mutagenesis techniques.

Such variants include those produced by nucleotide substitutions, deletions or additions, which may involve one or more nucleotides. The variants may be altered in coding regions, non-coding regions, or both. Alterations in the coding regions may produce conservative or non-conservative amino acid substitutions, deletions or additions. Especially preferred among these are silent substitutions, additions and deletions, which do not alter the properties and activities of the I-FLICE-1 or I-FLICE-2 protein or portions thereof. Also especially preferred in this regard are conservative substitutions.

Further embodiments of the invention include isolated nucleic acid molecules comprising a polynucleotide having a nucleotide sequence at least 95%, 96%, 97%, 98% or 99% identical to (a) a nucleotide sequence encoding the polypeptide having the amino acid sequence in SEQ ID NO:2; (b) a nucleotide sequence encoding the polypeptide having the amino acid sequence in SEQ ID NO:2, but lacking the N-terminal methionine; (c) a nucleotide sequence encoding the polypeptide having the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. 209038; or (d) a nucleotide sequence complementary to any of the nucleotide sequences in (a), (b), or (c).

Further embodiments of the invention include isolated nucleic acid molecules comprising a polynucleotide having a nucleotide sequence at least 95%, 96%, 97%, 98% or 99% identical to (a) a nucleotide sequence encoding the polypeptide having the amino acid sequence in SEQ ID NO:6; (b) a nucleotide sequence encoding the polypeptide having the amino acid sequence in SEQ ID NO:6, but lacking the N-terminal methionine; (c) a nucleotide sequence encoding the polypeptide having the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. 209041; or (d) a nucleotide sequence complementary to any of the nucleotide sequences in (a), (b), or (c).

Additional embodiments of the invention include isolated nucleic acid molecules comprising a polynucleotide having a nucleotide sequence at least

95%, 96%, 97%, 98% or 99% identical to (a) a nucleotide sequence encoding a polypeptide comprising amino acids from about 1 to about 75 in SEQ ID NO:2; (b) a nucleotide sequence encoding a polypeptide comprising amino acids from about 91 to about 171 in SEQ ID NO:2; (c) a nucleotide sequence encoding a polypeptide comprising amino acids from about 172 to about 375 in SEQ ID NO:2; (d) a nucleotide sequence encoding a polypeptide comprising amino acids from about 376 to about 480 in SEQ ID NO:2; (e) a nucleotide sequence encoding a polypeptide comprising amino acids from about 1 to about 75 in SEQ ID NO:6; (f) a nucleotide sequence encoding a polypeptide comprising amino acids from about 76 to about 252 in SEQ ID NO:6; (g) a nucleotide sequence encoding a polypeptide comprising amino acids from about 253 to about 348 in SEQ ID NO:6; (h) or a nucleotide sequence complementary to any of the nucleotide sequences in (a), (b), (c), (d), (e), (f), or (g).

By a polynucleotide having a nucleotide sequence at least, for example, 95% "identical" to a reference nucleotide sequence encoding an I-FLICE polypeptide is intended that the nucleotide sequence of the polynucleotide is identical to the reference sequence except that the polynucleotide sequence may include up to five point mutations per each 100 nucleotides of the reference nucleotide sequence encoding an I-FLICE polypeptide. In other words, to obtain a polynucleotide having a nucleotide sequence at least 95% identical to a reference nucleotide sequence, up to 5% of the nucleotides in the reference sequence may be deleted or substituted with another nucleotide, or a number of nucleotides up to 5% of the total nucleotides in the reference sequence may be inserted into the reference sequence. These mutations of the reference sequence may occur at the 5' or 3' terminal positions of the reference nucleotide sequence or anywhere between those terminal positions, interspersed either individually among nucleotides in the reference sequence or in one or more contiguous groups within the reference sequence.

As a practical matter, whether any particular nucleic acid molecule is at least 95%, 96%, 97%, 98% or 99% identical to, for instance, the nucleotide

sequence shown in SEQ ID NO:1 or SEQ ID NO:5 or to the nucleotides sequence of the deposited cDNA clone can be determined conventionally using known computer programs such as the Bestfit program (Wisconsin Sequence Analysis Package, Version 8 for Unix, Genetics Computer Group, University Research Park, 575 Science Drive, Madison, WI 53711). Bestfit uses the local homology algorithm of Smith and Waterman, *Advances in Applied Mathematics* 2:482-489 (1981), to find the best segment of homology between two sequences. When using Bestfit or any other sequence alignment program to determine whether a particular sequence is, for instance, 95% identical to a reference sequence according to the present invention, the parameters are set, of course, such that the percentage of identity is calculated over the full length of the reference nucleotide sequence and that gaps in homology of up to 5% of the total number of nucleotides in the reference sequence are allowed.

The present application is directed to nucleic acid molecules at least 95%, 96%, 97%, 98% or 99% identical to the nucleic acid sequence shown in SEQ ID NO:1 or SEQ ID NO:5 or to the nucleic acid sequence of the deposited cDNAs, irrespective of whether they encode a polypeptide having I-FLICE activity. This is because even where a particular nucleic acid molecule does not encode a polypeptide having I-FLICE activity, one of skill in the art would still know how to use the nucleic acid molecule, for instance, as a hybridization probe or a polymerase chain reaction (PCR) primer. Uses of the nucleic acid molecules of the present invention that do not encode a polypeptide having I-FLICE activity include, *inter alia*, (1) isolating the I-FLICE-1 or I-FLICE-2 gene or allelic variants thereof in a cDNA library; (2) *in situ* hybridization (e.g., "FISH") to metaphase chromosomal spreads to provide precise chromosomal location of the I-FLICE-1 or I-FLICE-2 gene, as described in Verma *et al.*, *Human Chromosomes: A Manual of Basic Techniques*, Pergamon Press, New York (1988); and (3) Northern Blot analysis for detecting I-FLICE-1 or I-FLICE-2 mRNA expression in specific tissues.

Preferred, however, are nucleic acid molecules having sequences at least 95%, 96%, 97%, 98% or 99% identical to a nucleic acid sequence shown in SEQ ID NO:1 or SEQ ID NO:5 or to a nucleic acid sequence of the deposited cDNA which do, in fact, encode a polypeptide having I-FLICE protein activity. By "a polypeptide having I-FLICE activity" is intended polypeptides exhibiting I-FLICE-1 or I-FLICE-2 activity in a particular biological assay. For example, I-FLICE-1 or I-FLICE-2 protein activity can be measured using the cell death assay as described in Example 6.

Of course, due to the degeneracy of the genetic code, one of ordinary skill in the art will immediately recognize that a large number of the nucleic acid molecules having a sequence at least 95%, 96%, 97%, 98%, or 99% identical to a nucleic acid sequence of the deposited cDNA or a nucleic acid sequence shown in SEQ ID NO:1 or SEQ ID NO:5 will encode "a polypeptide having I-FLICE activity." In fact, since degenerate variants of these nucleotide sequences all encode the same polypeptide, this will be clear to the skilled artisan even without performing the above described comparison assay. It will be further recognized in the art that, for such nucleic acid molecules that are not degenerate variants, a reasonable number will also encode a polypeptide having I-FLICE activity. This is because the skilled artisan is fully aware of amino acid substitutions that are either less likely or not likely to significantly effect protein function (e.g., replacing one aliphatic amino acid with a second aliphatic amino acid).

For example, guidance concerning how to make phenotypically silent amino acid substitutions is provided in Bowie, J. U. *et al.*, "Deciphering the Message in Protein Sequences: Tolerance to Amino Acid Substitutions," *Science* 247:1306-1310 (1990), wherein the authors indicate that proteins are surprisingly tolerant of amino acid substitutions.



### *Vectors and Host Cells*

The present invention also relates to vectors which include the isolated DNA molecules of the present invention, host cells which are genetically engineered with the recombinant vectors, and the production of I-FLICE-1 or I-FLICE-2 polypeptides or fragments thereof by recombinant techniques.

The polynucleotides may be joined to a vector containing a selectable marker for propagation in a host. Generally, a plasmid vector is introduced in a precipitate, such as a calcium phosphate precipitate, or in a complex with a charged lipid. If the vector is a virus, it may be packaged *in vitro* using an appropriate packaging cell line and then transduced into host cells.

The DNA insert should be operatively linked to an appropriate promoter, such as the phage lambda PL promoter, the *E. coli lac*, *trp* and *tac* promoters, the SV40 early and late promoters and promoters of retroviral LTRs, to name a few. Other suitable promoters will be known to the skilled artisan. The expression constructs will further contain sites for transcription initiation, termination and, in the transcribed region, a ribosome binding site for translation. The coding portion of the mature transcripts expressed by the constructs will preferably include a translation initiating at the beginning and a termination codon (UAA, UGA or UAG) appropriately positioned at the end of the polypeptide to be translated.

As indicated, the expression vectors will preferably include at least one selectable marker. Such markers include dihydrofolate reductase or neomycin resistance for eukaryotic cell culture and tetracycline or ampicillin resistance genes for culturing in *E. coli* and other bacteria. Representative examples of appropriate hosts include, but are not limited to, bacterial cells, such as *E. coli*, *Streptomyces* and *Salmonella typhimurium* cells; fungal cells, such as yeast cells; insect cells such as *Drosophila* S2 and *Spodoptera* Sf9 cells; animal cells such as CHO, COS and Bowes melanoma cells; and plant cells. Appropriate culture mediums and conditions for the above-described host cells are known in the art.

Among vectors preferred for use in bacteria include pQE70, pQE60 and pQE-9, available from Qiagen; pBS vectors, Phagescript vectors, Bluescript vectors, pNH8A, pNH16a, pNH18A, pNH46A, available from Stratagene; and ptrc99a, pKK223-3, pKK233-3, pDR540, pRIT5 available from Pharmacia. Among preferred eukaryotic vectors are pWLNEO, pSV2CAT, pOG44, pXT1 and pSG available from Stratagene; and pSVK3, pBPV, pMSG and pSVL available from Pharmacia. Other suitable vectors will be readily apparent to the skilled artisan.

Introduction of the construct into the host cell can be effected by calcium phosphate transfection, DEAE-dextran mediated transfection, cationic lipid-mediated transfection, electroporation, transduction, infection or other methods. Such methods are described in many standard laboratory manuals, such as Davis *et al.*, *Basic Methods In Molecular Biology* (1986).

The polypeptide may be expressed in a modified form, such as a fusion protein, and may include not only secretion signals, but also additional heterologous functional regions. For instance, a region of additional amino acids, particularly charged amino acids, may be added to the N-terminus of the polypeptide to improve stability and persistence in the host cell, during purification, or during subsequent handling and storage. Also, peptide moieties may be added to the polypeptide to facilitate purification. Such regions may be removed prior to final preparation of the polypeptide. The addition of peptide moieties to polypeptides to engender secretion or excretion, to improve stability and to facilitate purification, among others, are familiar and routine techniques in the art. A preferred fusion protein comprises a heterologous region from immunoglobulin that is useful to solubilize proteins. For example, EP-A-O 464 533 (Canadian counterpart 2045869) discloses fusion proteins comprising various portions of constant region of immunoglobulin molecules together with another human protein or part thereof. In many cases, the Fc part in a fusion protein is thoroughly advantageous for use in therapy and diagnosis and thus results, for example, in improved pharmacokinetic properties (EP-A 0232 262). On the other

hand, for some uses it would be desirable to be able to delete the Fc part after the fusion protein has been expressed, detected and purified in the advantageous manner described. This is the case when Fc portion proves to be a hindrance to use in therapy and diagnosis, for example when the fusion protein is to be used as antigen for immunizations. In drug discovery, for example, human proteins, such as, hIL5-receptor has been fused with Fc portions for the purpose of high-throughput screening assays to identify antagonists of hIL-5. See, D. Bennett *et al.*, *Journal of Molecular Recognition*, Vol. 8:52-58 (1995) and K. Johanson *et al.*, *The Journal of Biological Chemistry*, Vol. 270, No. 16:9459-9471 (1995).

The I-FLICE-1 or I-FLICE-2 protein can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, high performance liquid chromatography ("HPLC") is employed for purification. Polypeptides of the present invention include naturally purified products, products of chemical synthetic procedures, and products produced by recombinant techniques from a prokaryotic or eukaryotic host, including, for example, bacterial, yeast, higher plant, insect and mammalian cells. Depending upon the host employed in a recombinant production procedure, the polypeptides of the present invention may be glycosylated or may be non-glycosylated. In addition, polypeptides of the invention may also include an initial modified methionine residue, in some cases as a result of host-mediated processes.

#### ***I-FLICE-1 and I-FLICE-2 Polypeptides and Fragments***

The invention further provides an isolated I-FLICE-1 or I-FLICE-2 polypeptide having the amino acid sequence encoded by the deposited cDNAs,

or the amino acid sequence in SEQ ID NO:2 or SEQ ID NO:6, or a peptide or polypeptide comprising a portion of the above polypeptides.

It will be recognized in the art that some amino acid sequences of the I-FLICE-1 or I-FLICE-2 polypeptide can be varied without significant effect of the structure or function of the protein. If such differences in sequence are contemplated, it should be remembered that there will be critical areas on the protein which determine activity.

Thus, the invention further includes variations of the I-FLICE-1 or I-FLICE-2 polypeptide which show substantial I-FLICE-1 or I-FLICE-2 polypeptide activity or which include regions of I-FLICE-1 or I-FLICE-2 protein such as the protein portions discussed below. Such mutants include deletions, insertions, inversions, repeats, and type substitutions. As indicated above, guidance concerning which amino acid changes are likely to be phenotypically silent can be found in Bowie, J.U., *et al.*, "Deciphering the Message in Protein Sequences: Tolerance to Amino Acid Substitutions," *Science* 247:1306-1310 (1990).

Thus, the fragment, derivative or analog of the polypeptide of SEQ ID NO:2 or SEQ ID NO:6, or that encoded by the deposited cDNAs, may be (i) one in which one or more of the amino acid residues are substituted with a conserved or non-conserved amino acid residue (preferably a conserved amino acid residue) and such substituted amino acid residue may or may not be one encoded by the genetic code, or (ii) one in which one or more of the amino acid residues includes a substituent group, or (iii) one in which the mature polypeptide is fused with another compound, such as a compound to increase the half-life of the polypeptide (for example, polyethylene glycol), or (iv) one in which the additional amino acids are fused to the mature polypeptide, such as an IgG Fc fusion region peptide or leader or secretory sequence or a sequence which is employed for purification of the mature polypeptide or a proprotein sequence. Such fragments, derivatives and analogs are deemed to be within the scope of those skilled in the art from the teachings herein.

Of particular interest are substitutions of charged amino acids with another charged amino acid and with neutral or negatively charged amino acids. The latter results in proteins with reduced positive charge to improve the characteristics of the I-FLICE-1 or I-FLICE-2 protein. The prevention of aggregation is highly desirable. Aggregation of proteins not only results in a loss of activity but can also be problematic when preparing pharmaceutical formulations, because they can be immunogenic. (Pinckard *et al.*, *Clin. Exp. Immunol.* 2:331-340 (1967); Robbins *et al.*, *Diabetes* 36:838-845 (1987); Cleland *et al.* *Crit. Rev. Therapeutic Drug Carrier Systems* 10:307-377 (1993)).

The replacement of amino acids can also change the selectivity of binding to cell surface receptors. Ostade *et al.*, *Nature* 361:266-268 (1993) describes certain mutations resulting in selective binding of TNF- $\alpha$  to only one of the two known types of TNF receptors. Thus, the I-FLICE-1 or I-FLICE-2 of the present invention may include one or more amino acid substitutions, deletions or additions, either from natural mutations or human manipulation.

As indicated, changes are preferably of a minor nature, such as conservative amino acid substitutions that do not significantly affect the folding or activity of the protein (see Table 1).

TABLE 1. Conservative Amino Acid Substitutions.

Aromatic	Phenylalanine Tryptophan Tyrosine
Hydrophobic	Leucine Isoleucine Valine
Polar	Glutamine Asparagine
Basic	Arginine Lysine Histidine
Acidic	Aspartic Acid Glutamic Acid
Small	Alanine Serine Threonine Methionine Glycine

Of course, the number of amino acid substitutions a skilled artisan would make depends on many factors, including those described above and below. Generally speaking, the number of substitutions for any given I-FLICE-1 or I-FLICE-2 polypeptide, or mutant thereof, will not be more than 50, 40, 30, 20, 10, 5, or 3, depending on the objective.

Amino acids in the I-FLICE-1 or I-FLICE-2 protein of the present invention that are essential for function can be identified by methods known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (Cunningham and Wells, *Science* 244:1081-1085 (1989)). The latter procedure introduces single alanine mutations at every residue in the molecule. Sites that are critical for ligand interactions can also be determined by structural analysis such as crystallization, nuclear magnetic resonance or photoaffinity labeling (Smith *et al.*, *J. Mol. Biol.* 224:899-904 (1992) and de Vos *et al.* *Science* 255:306-312 (1992)).

The polypeptides of the present invention are preferably provided in an isolated form. By "isolated polypeptide" is intended a polypeptide removed from its native environment. Thus, a polypeptide produced and/or contained within a recombinant host cell is considered isolated for purposes of the present invention. Also intended as an "isolated polypeptide" are polypeptides that have been purified, partially or substantially, from a recombinant host cell. For example, a recombinantly produced version of the I-FLICE-1 or I-FLICE-2 polypeptide can be substantially purified by the one-step method described in Smith and Johnson, *Gene* 67:31-40 (1988).

The polypeptides of the present invention include the polypeptide encoded by the deposited cDNA; a polypeptide comprising amino acids about 1 to about 480 in SEQ ID NO:2; a polypeptide comprising amino acids about 2 to about 480 in SEQ ID NO:2; as well as polypeptides which are at least 95% identical, still more preferably at least 96%, 97%, 98% or 99% identical to those described above and also include portions of such polypeptides with at least 30 amino acids and more preferably at least 50 amino acids.

The polypeptides of the present invention also include the polypeptide encoded by the deposited cDNA; a polypeptide comprising amino acids about 1 to about 348 in SEQ ID NO:6; a polypeptide comprising amino acids about 2 to about 348 in SEQ ID NO:6; as well as polypeptides which are at least 95% identical, still more preferably at least 96%, 97%, 98% or 99% identical to those described above and also include portions of such polypeptides with at least 30 amino acids and more preferably at least 50 amino acids.

The polypeptides of the present invention further include the polypeptide comprising amino acids from about 1 to about 75 in SEQ ID NO:2; amino acids from about 91 to about 171 in SEQ ID NO:2; amino acids from about 172 to about 375 in SEQ ID NO:2; amino acids from about 376 to about 480 in SEQ ID NO:2; amino acids from about 1 to about 75 in SEQ ID NO:6; amino acids from about 76 to about 252 in SEQ ID NO:6; amino acids from about 253 to about 348 in SEQ ID NO:6; as well as polypeptides which are at least 95% identical, still

more preferably at least 96%, 97%, 98% or 99% identical to those described above and also include portions of such polypeptides with at least 30 amino acids and more preferably at least 50 amino acids.

By a polypeptide having an amino acid sequence at least, for example, 95% "identical" to a reference amino acid sequence of an I-FLICE-1 or I-FLICE-2 polypeptide is intended that the amino acid sequence of the polypeptide is identical to the reference sequence except that the polypeptide sequence may include up to five amino acid alterations per each 100 amino acids of the reference amino acid of the I-FLICE-1 or I-FLICE-2 polypeptide. In other words, to obtain a polypeptide having an amino acid sequence at least 95% identical to a reference amino acid sequence, up to 5% of the amino acid residues in the reference sequence may be deleted or substituted with another amino acid, or a number of amino acids up to 5% of the total amino acid residues in the reference sequence may be inserted into the reference sequence. These alterations of the reference sequence may occur at the amino or carboxy terminal positions of the reference amino acid sequence or anywhere between those terminal positions, interspersed either individually among residues in the reference sequence or in one or more contiguous groups within the reference sequence.

As a practical matter, whether any particular polypeptide is at least 95%, 96%, 97%, 98% or 99% identical to, for instance, the amino acid sequence shown in SEQ ID NO:2 or SEQ ID NO:6 or to the amino acid sequence encoded by deposited cDNA clones can be determined conventionally using known computer programs such the Bestfit program (Wisconsin Sequence Analysis Package, Version 8 for Unix, Genetics Computer Group, University Research Park, 575 Science Drive, Madison, WI 53711). When using Bestfit or any other sequence alignment program to determine whether a particular sequence is, for instance, 95% identical to a reference sequence according to the present invention, the parameters are set, of course, such that the percentage of identity is calculated over the full length of the reference amino acid sequence and that gaps in



-27-

homology of up to 5% of the total number of amino acid residues in the reference sequence are allowed.

The polypeptide of the present invention are useful as a molecular weight marker on SDS-PAGE gels or on molecular sieve gel filtration columns using methods well known to those of skill in the art.

In another aspect, the invention provides a peptide or polypeptide comprising an epitope-bearing portion of a polypeptide of the invention. The epitope of this polypeptide portion is an immunogenic or antigenic epitope of a polypeptide described herein. An "immunogenic epitope" is defined as a part of a protein that elicits an antibody response when the whole protein is the immunogen. On the other hand, a region of a protein molecule to which an antibody can bind is defined as an "antigenic epitope." The number of immunogenic epitopes of a protein generally is less than the number of antigenic epitopes. See, for instance, Geysen *et al.*, *Proc. Natl. Acad. Sci. USA* 81:3998-4002 (1983).

As to the selection of peptides or polypeptides bearing an antigenic epitope (i.e., that contain a region of a protein molecule to which an antibody can bind), it is well known in that art that relatively short synthetic peptides that mimic part of a protein sequence are routinely capable of eliciting an antiserum that reacts with the partially mimicked protein. See, for instance, Sutcliffe, J. G., Shinnick, T. M., Green, N. and Learner, R.A. (1983) Antibodies that react with predetermined sites on proteins. *Science* 219:660-666. Peptides capable of eliciting protein-reactive sera are frequently represented in the primary sequence of a protein, can be characterized by a set of simple chemical rules, and are confined neither to immunodominant regions of intact proteins (i.e., immunogenic epitopes) nor to the amino or carboxyl terminals.

Antigenic epitope-bearing peptides and polypeptides of the invention are therefore useful to raise antibodies, including monoclonal antibodies, that bind specifically to a polypeptide of the invention. See, for instance, Wilson *et al.*, *Cell* 37:767-778 (1984) at 777.

Antigenic epitope-bearing peptides and polypeptides of the invention preferably contain a sequence of at least seven, more preferably at least nine and most preferably between about at least about 15 to about 30 amino acids contained within the amino acid sequence of a polypeptide of the invention.

5 Non-limiting examples of antigenic polypeptides or peptides that can be used to generate I-FLICE-1 -specific antibodies include: a polypeptide comprising amino acid residues from about 41 to about 92 in SEQ ID NO:2; a polypeptide comprising amino acid residues from about 155 to about 249 in SEQ ID NO:2; a polypeptide comprising amino acid residues from about 332 to about 474 in  
10 SEQ ID NO:2. As indicated above, the inventors have determined that the above polypeptide fragments are antigenic regions of the I-FLICE-1 protein.

Non-limiting examples of antigenic polypeptides or peptides that can be used to generate I-FLICE-2 -specific antibodies include: a polypeptide comprising amino acid residues from about 62 to about 136 in SEQ ID NO:6; a polypeptide  
15 comprising amino acid residues from about 184 to about 193 in SEQ ID NO:6; a polypeptide comprising amino acid residues from about 205 to about 341 in SEQ ID NO:6. The inventors have determined that the above polypeptide fragments are antigenic regions of the I-FLICE-2 protein.

The epitope-bearing peptides and polypeptides of the invention may be  
20 produced by any conventional means. Houghten, R. A. (1985) General method for the rapid solid-phase synthesis of large numbers of peptides: specificity of antigen-antibody interaction at the level of individual amino acids. *Proc. Natl. Acad. Sci. USA* 82:5131-5135. This "Simultaneous Multiple Peptide Synthesis (SMPS)" process is further described in U.S. Patent No. 4,631,211 to Houghten  
25 *et al.* (1986).

As one of skill in the art will appreciate, I-FLICE-1 or I-FLICE-2 polypeptides of the present invention and the epitope-bearing fragments thereof described above can be combined with parts of the constant domain of immunoglobulins (IgG), resulting in chimeric polypeptides. These fusion  
30 proteins facilitate purification and show an increased half-life *in vivo*. This has

been shown, e.g., for chimeric proteins consisting of the first two domains of the human CD4-polypeptide and various domains of the constant regions of the heavy or light chains of mammalian immunoglobulins (EPA 394,827; Traunecker *et al.*, *Nature* 331:84- 86 (1988)). Fusion proteins that have a disulfide-linked dimeric structure due to the IgG part can also be more efficient in binding and neutralizing other molecules than the monomeric I-FLICE-1 or I-FLICE-2 protein or protein fragment alone (Fountoulakis *et al.*, *J. Biochem* 270:3958-3964 (1995)).

### ***Disease Diagnosis and Prognosis***

It is believed that certain tissues in mammals with specific disease states associated with aberrant cell survival express significantly altered levels of I-FLICE-1 or I-FLICE-2 and mRNA encoding I-FLICE-1 or I-FLICE-2 when compared to a corresponding "standard" mammal, i.e., a mammal of the same species not having the disease state. Thus, the present invention is useful for detecting such states in mammals. Preferred mammals include monkeys, apes, cats, dogs, cows, pigs, horses, rabbits and humans. Particularly preferred are humans.

It is further believed that enhanced levels of I-FLICE-1 or I-FLICE-2 can be detected in certain body fluids (e.g., sera, plasma, urine, and spinal fluid) from mammals with the disease state when compared to analogous fluids from mammals of the same species not having the disease state. Thus, the invention provides a diagnostic method useful during diagnosis of disease states, which involves assaying the expression level of the gene encoding I-FLICE-1 or I-FLICE-2 in mammalian cells or body fluid and comparing the gene expression level with a standard I-FLICE-1 or I-FLICE-2, whereby an increase or decrease in the gene expression level over the standard is indicative of certain disease states associated with aberrant cell survival.

Where diagnosis of a disease state involving I-FLICE-1 or I-FLICE-2 of the present invention has already been made according to conventional methods, the present invention is useful as a prognostic indicator, whereby patients exhibiting significantly aberrant I-FLICE-1 or I-FLICE-2 gene expression levels will experience a worse clinical outcome relative to patients expressing the gene at a lower level.

By "assaying the expression level of the gene encoding I-FLICE-1 or I-FLICE-2" is intended qualitatively or quantitatively measuring or estimating the level of I-FLICE-1 or I-FLICE-2 protein or the level of the mRNA encoding I-FLICE-1 or I-FLICE-2 protein in a first biological sample either directly (e.g., by determining or estimating absolute protein level or mRNA level) or relatively (e.g., by comparing to the I-FLICE-1 or I-FLICE-2 protein level or mRNA level in a second biological sample).

Preferably, the I-FLICE-1 or I-FLICE-2 protein level or mRNA level in the first biological sample is measured or estimated and compared to a standard I-FLICE-1 or I-FLICE-2 protein level or mRNA level, the standard being taken from a second biological sample obtained from an individual not having the disease state. As will be appreciated in the art, once a standard I-FLICE-1 or I-FLICE-2 protein level or mRNA level is known, it can be used repeatedly as a standard for comparison.

By "biological sample" is intended any biological sample obtained from an individual, cell line, tissue culture, or other source which contains I-FLICE-1 or I-FLICE-2 protein or mRNA. Biological samples include mammalian body fluids (such as sera, plasma, urine, synovial fluid and spinal fluid) which contain I-FLICE-1 or I-FLICE-2 protein, and ovarian, prostate, heart, placenta, pancreas liver, spleen, lung, breast, umbilical tissue, and other tissues. Methods for obtaining tissue biopsies and body fluids from mammals are well known in the art. Where the biological sample is to include mRNA, a tissue biopsy is the preferred source.

-31-

Diseases associated with increased cell survival, or the inhibition of apoptosis, include cancers (such as follicular lymphomas, carcinomas with p53 mutations, hormone-dependent tumors, and cancers of the breast, ovary, prostate, bone, liver, lung, pancreas, and spleen); autoimmune disorders (such as systemic lupus erythematosus and immune-related glomerulonephritis rheumatoid arthritis) and viral infections (such as herpes viruses, pox viruses and adenoviruses), information graft v. host disease, acute graft rejection, and chronic graft rejection. Diseases associated with decreased cell survival, or increased apoptosis, include Alzheimer's disease, Parkinson's disease, rheumatoid arthritis, septic shock, sepsis, stroke, CNS inflammation, osteoporosis, ischemia, reperfusion injury, cell death associated with cardiovascular disease, polycystic kidney disease, apoptosis of endothelial cells in cardiovascular disease, degenerative liver disease, MS and head injury damage.

Assays available to detect levels of proteins are well known to those of skill in the art, for example, radioimmunoassays, competitive-binding assays, Western blot analysis, and preferably an ELISA assay may be employed.

I-FLICE-1 or I-FLICE-2 specific antibodies can be raised against the intact I-FLICE-1 or I-FLICE-2 protein or an antigenic polypeptide fragment thereof, which may presented together with a carrier protein, such as an albumin, to an animal system (such as rabbit or mouse) or, if it is long enough (at least about 25 amino acids), without a carrier.

As used herein, the term "antibody" (Ab) or "monoclonal antibody" (mAb) is meant to include intact molecules as well as antibody fragments (such as, for example, Fab and F(ab')<sub>2</sub> fragments) which are capable of specifically binding to the I-FLICE-1 or I-FLICE-2 protein. Fab and F(ab')<sub>2</sub> fragments lack the Fc fragment of intact antibody, clear more rapidly from the circulation, and may have less non-specific tissue binding of an intact antibody (Wahl *et al.*, *J. Nucl. Med.* 24:316-325 (1983)). Thus, these fragments are preferred.

The antibodies of the present invention may be prepared by any of a variety of methods. For example, cells expressing the I-FLICE-1 or I-FLICE-2

protein or an antigenic fragment thereof can be administered to an animal in order to induce the production of sera containing polyclonal antibodies. In a preferred method, a preparation of I-FLICE-1 or I-FLICE-2 protein is prepared and purified to render it substantially free of natural contaminants. Such a preparation is then introduced into an animal in order to produce polyclonal antisera of greater specific activity.

In the most preferred method, the antibodies of the present invention are monoclonal antibodies (or I-FLICE-1 or I-FLICE-2 protein binding fragments thereof). Such monoclonal antibodies can be prepared using hybridoma technology (Kohler *et al.*, *Nature* 256:495 (1975); Kohler *et al.*, *Eur. J. Immunol.* 6:511 (1976); Kohler *et al.*, *Eur. J. Immunol.* 6:292 (1976); Hammerling *et al.*, In: *Monoclonal Antibodies and T-Cell Hybridomas*, Elsevier, N.Y., (1981) pp. 563-681).

Assaying I-FLICE-1 or I-FLICE-2 protein levels in a biological sample can occur using antibody-based techniques. For example, I-FLICE-1 or I-FLICE-2 protein expression in tissues can be studied with classical immunohistological methods (Jalkanen, M., *et al.*, *J. Cell. Biol.* 101:976-985 (1985); Jalkanen, M., *et al.*, *J. Cell. Biol.* 105:3087-3096 (1987)).

As noted above, other antibody-based methods useful for detecting I-FLICE-1 or I-FLICE-2 protein gene expression include immunoassays, such as the enzyme linked immunosorbent assay (ELISA) and the radioimmunoassay (RIA).

Suitable labels are known in the art and include enzyme labels, such as, Glucose oxidase, and radioisotopes, such as iodine ( $^{125}\text{I}$ ,  $^{121}\text{I}$ ), carbon ( $^{14}\text{C}$ ), sulfur ( $^{35}\text{S}$ ), tritium ( $^3\text{H}$ ), indium ( $^{112}\text{In}$ ), and technetium ( $^{99\text{m}}\text{Tc}$ ), and fluorescent labels, such as fluorescein and rhodamine, and biotin.

Total cellular RNA can be isolated from a biological sample using the single-step guanidinium-thiocyanate-phenol-chloroform method described in Chomczynski and Sacchi, *Anal. Biochem.* 162:156-159 (1987). Levels of mRNA encoding the I-FLICE-1 or I-FLICE-2 protein are then assayed using any

appropriate method. These include Northern blot analysis (Harada *et al.*, *Cell* 63:303-312 (1990)), S1 nuclease mapping (Fujita *et al.*, *Cell* 49:357- 367 (1987)), the polymerase chain reaction (PCR), reverse transcription in combination with the polymerase chain reaction (RT-PCR) (Makino *et al.*,  
5 *Technique* 2:295-301 (1990)), and reverse transcription in combination with the ligase chain reaction (RT-LCR).

### ***Agonists and Antagonists - Assays and Molecules***

The invention also provides a method of screening compounds to identify  
10 agonists and antagonists of I-FLICE-1 or I-FLICE-2. By "agonist" is intended naturally occurring and synthetic compounds capable of enhancing one or more activity mediated by I-FLICE-1 or I-FLICE-2 polypeptides. By "antagonist" is intended naturally occurring and synthetic compounds capable of inhibiting one or more activity mediated by I-FLICE-1 or I-FLICE-2 polypeptides.

Thus, in a further aspect, a screening method is provided for determining  
15 whether a candidate agonist or antagonist is capable of enhancing or inhibiting a cellular activity of either an I-FLICE-1 or I-FLICE-2 polypeptide. The method involves contacting cells which express one or both of the I-FLICE-1 or I-FLICE-2 polypeptides with a candidate compound, assaying a cellular response, and comparing the cellular response to a standard cellular response, the standard being  
20 assayed in absence of the candidate compound, whereby an increased cellular response over the standard indicates that the candidate compound is an agonist of the polypeptide activity and a decreased cellular response compared to the standard indicates that the candidate compound is an antagonist of the activity. By "assaying a cellular response" is intended qualitatively or quantitatively  
25 measuring a cellular response in the presence of a candidate compound and either an I-FLICE-1 or I-FLICE-2 polypeptide (e.g., decreased or increased TNFR-1 or CD-95 induced apoptosis, binding of I-FLICE-1 or I-FLICE-2 to natural cellular ligands such as FLICE and Mch4/FLICE2).

Potential antagonists include small organic molecules amino acid sequences which bind to I-FLICE-1 or I-FLICE-2, fragments of I-FLICE-1 and I-FLICE-2, as well as anti-I-FLICE-1 and anti-I-FLICE-2 antibodies. Fragments of I-FLICE-1 and I-FLICE-2, which may be naturally occurring or synthetic, antagonize I-FLICE-1 and I-FLICE-2 polypeptide mediated activities by competing for binding to natural cellular ligands. Small organic molecules can antagonize I-FLICE-1 and I-FLICE-2 polypeptide mediated activities by binding either competitively or non-competitively to I-FLICE-1 or I-FLICE-2 or a cellular ligand of these proteins. Examples of small molecules include but are not limited to nucleotide sequences and small peptides or peptide-like molecules. Such molecules may be produced and screened for activity by a variety of methods (e.g., Light and Lerner, *Bioorganic & Medicinal Chemistry* 3(7):955-967 (1995); Cheng *et al.*, *Gene* 171:1-8 (1996); Gates *et al.*, *J. Mol. Biol.* 255:373-386 (1996)).

Similarly, potential agonists also include fragments of the polypeptides of the present invention, as well as anti-I-FLICE-1 and anti-I-FLICE-2 antibodies. Fragments of these proteins can act as agonists of I-FLICE-1 and I-FLICE-2 polypeptide mediated activities by binding to natural cellular ligands and inducing activities associated with the full-length protein. Agonists and antagonists of the present invention also include amino acid sequences having 95% or more identity to those shown in SEQ ID NOs:2 and 6, or fragments thereof.

Other potential antagonists include antisense oligonucleotides and oligonucleotides capable of forming triple helices with the sequences shown in SEQ ID NOs:1 and 5. Once a gene sequence is known, antisense and triple helix technologies can be used to regulate gene expression. Okano, *J. Neurochem.* 56:560 (1991); OLIGONUCLEOTIDES AS INHIBITORS OF GENE EXPRESSION, CRC Press, Boca Raton, FL. (1988); Dervan *et al.*, *Science* 251:1360 (1991); Cooney *et al.*, *Science* 241:456 (1988); Lee *et al.*, *Nucl. Acids Res.* 6:3073 (1979). In regards to antisense technology, for example, an oligonucleotide may be designed which is complementary to a portion of the I-FLICE-1 or I-FLICE-2 DNA



sequences which is transcribed into RNA. This oligonucleotide may be delivered to cells in a number of forms, including as antisense RNA or incorporated into an expression vector. If incorporated into an expression vector, the oligonucleotide is generally orientated in a manner that an RNA molecule is produced upon *in vivo* expression which is complementary to that of the I-FLICE-1 or I-FLICE-2 mRNA sequence. The expressed antisense RNA molecule will hybridize to I-FLICE-1 or I-FLICE-2 mRNA and block translation *in vivo*.

The experiments set forth in Example 5 demonstrate that I-FLICE-1 binds to both FLICE and Mch4/FLICE2. Immunoprecipitation assays similar to that described in Example 5 can be used to identify additional molecules which bind to I-FLICE-1 and I-FLICE-2. Such binding molecules are candidate antagonists and agonists.

Example 6 sets forth a cell death assay used to demonstrate that overexpression of I-FLICE-1 results in the inhibition of TNFR-1 and CD-95 induced cell death. This assay can also be used to screen for compounds having agonistic and antagonistic activity directed to I-FLICE-1 and I-FLICE-2. Such a screening method is used to determine whether the compound increases or decreases TNFR-1 and CD-95 induced cell death in the presence of I-FLICE-1 or I-FLICE-2 either individually or in combination.

Proteins and other compounds which bind the I-FLICE-1 or I-FLICE-2 polypeptide domains are also candidate agonists and antagonists according to the present invention. Such binding compounds can be "captured" using the yeast two-hybrid system (Fields and Song, *Nature* 340:245-246 (1989); Gyuris *et al.*, *Cell* 75:791-803 (1993); Zervos *et al.*, *Cell* 72:223 -232 (1993)).

The agonists may be employed for instance to enhance the action of I-FLICE-1 or I-FLICE-2 polypeptides, for example, in the treatment of Alzheimer's disease, Parkinson's disease, rheumatoid arthritis, septic shock, sepsis, stroke, CNS inflammation, osteoporosis, ischemia, reperfusion injury, cell death associated with cardiovascular disease, polycystic kidney disease, apoptosis of

endothelial cells in cardiovascular disease, degenerative liver disease, MS and head injury damage.

The antagonists may be employed for instance to inhibit the action of I-FLICE-1 or I-FLICE-2 polypeptides, for example, in the treatment of cancers (such as follicular lymphomas, carcinomas with p53 mutations, hormone-dependent tumors, and cancers of the breast, ovary, prostate, bone, liver, lung, pancreas, and spleen); autoimmune disorders (such as systemic lupus erythematosus and immune-related glomerulonephritis rheumatoid arthritis) and viral infections (such as herpes viruses, pox viruses and adenoviruses), information graft v. host disease, acute graft rejection, and chronic graft rejection.

The agonists and antagonists may be employed in a composition with a pharmaceutically acceptable carrier, e.g., as hereinafter described.

### *Therapeutics*

The novel mammalian inhibitors designated I-FLICE-1 and I-FLICE-2 (for inhibitor of FLICE) of the present invention, are catalytically inactive structural homologues of FLICE and Mch4/FLICE2 that inhibit both TNFR-1 and CD-95 induced apoptosis. These are the first examples of a naturally occurring catalytically inactive caspase that can act as a dominant negative inhibitor of apoptosis. The polypeptides of the present invention are useful for inhibiting apoptosis for therapeutic purposes, for example, in the treatment of Alzheimer's disease, Parkinson's disease, rheumatoid arthritis, septic shock, sepsis, stroke, CNS inflammation, osteoporosis, ischemia, reperfusion injury, cell death associated with cardiovascular disease, polycystic kidney disease, apoptosis of endothelial cells in cardiovascular disease, degenerative liver disease, MS and head injury damage.

### *Modes of Administration*

It will be appreciated that conditions caused by a decrease in the standard or normal level of I-FLICE-1 or I-FLICE-2 activity in an individual, can be treated by administration of I-FLICE-1 or I-FLICE-2 protein. Thus, the invention further provides a method of treating an individual in need of an increased level of I-FLICE-1 or I-FLICE-2 activity comprising administering to such an individual a pharmaceutical composition comprising an effective amount of an isolated I-FLICE-1 or I-FLICE-2 polypeptide of the invention, particularly a mature form of the I-FLICE-1 or I-FLICE-2, effective to increase the I-FLICE-1 or I-FLICE-2 activity level in such an individual.

As a general proposition, the total pharmaceutically effective amount of I-FLICE-1 or I-FLICE-2 polypeptide administered parenterally per dose will be in the range of about 1  $\mu\text{g/kg/day}$  to 10  $\text{mg/kg/day}$  of patient body weight, although, as noted above, this will be subject to therapeutic discretion. More preferably, this dose is at least 0.01  $\text{mg/kg/day}$ , and most preferably for humans between about 0.01 and 1  $\text{mg/kg/day}$  for the hormone. If given continuously, the I-FLICE-1 or I-FLICE-2 polypeptide is typically administered at a dose rate of about 1  $\mu\text{g/kg/hour}$  to about 50  $\mu\text{g/kg/hour}$ , either by 1-4 injections per day or by continuous subcutaneous infusions, for example, using a mini-pump. An intravenous bag solution may also be employed.

Pharmaceutical compositions containing the I-FLICE-1 or I-FLICE-2 of the invention may be administered orally, rectally, parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments, drops or transdermal patch), buccally, or as an oral or nasal spray. By "pharmaceutically acceptable carrier" is meant a non-toxic solid, semisolid or liquid filler, diluent, encapsulating material or formulation auxiliary of any type. The term "parenteral" as used herein refers to modes of administration which include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous and intraarticular injection and infusion.

### *Chromosome Assays*

The nucleic acid molecules of the present invention are also valuable for chromosome identification. The sequence is specifically targeted to and can hybridize with a particular location on an individual human chromosome. The mapping of DNAs to chromosomes according to the present invention is an important first step in correlating those sequences with genes associated with disease.

In certain preferred embodiments in this regard, the cDNA herein disclosed is used to clone genomic DNA of an I-FLICE-1 or I-FLICE-2 protein gene. This can be accomplished using a variety of well known techniques and libraries, which generally are available commercially. The genomic DNA then is used for *in situ* chromosome mapping using well known techniques for this purpose.

In addition, in some cases, sequences can be mapped to chromosomes by preparing PCR primers (preferably 15-25 bp) from the cDNA. Computer analysis of the 3' untranslated region of the gene is used to rapidly select primers that do not span more than one exon in the genomic DNA, thus complicating the amplification process. These primers are then used for PCR screening of somatic cell hybrids containing individual human chromosomes.

Fluorescence *in situ* hybridization ("FISH") of a cDNA clone to a metaphase chromosomal spread can be used to provide a precise chromosomal location in one step. This technique can be used with probes from the cDNA as short as 50 or 60 bp. For a review of this technique, see Verma *et al.*, *Human Chromosomes: A Manual Of Basic Techniques*, Pergamon Press, New York (1988).

Once a sequence has been mapped to a precise chromosomal location, the physical position of the sequence on the chromosome can be correlated with genetic map data. Such data are found, for example, in V. McKusick, *Mendelian Inheritance In Man*, available on-line through Johns Hopkins University, Welch

Medical Library. The relationship between genes and diseases that have been mapped to the same chromosomal region are then identified through linkage analysis (coinheritance of physically adjacent genes).

Next, it is necessary to determine the differences in the cDNA or genomic sequence between affected and unaffected individuals. If a mutation is observed in some or all of the affected individuals but not in any normal individuals, then the mutation is likely to be the causative agent of the disease.

Having generally described the invention, the same will be more readily understood by reference to the following examples, which are provided by way of illustration and are not intended as limiting.

### *Examples*

#### *Example 1(a): Expression and Purification of I-FLICE-1 in E. coli*

The bacterial expression vector pQE60 is used for bacterial expression in this example. (QIAGEN, Inc., 9259 Eton Avenue, Chatsworth, CA, 91311). pQE60 encodes ampicillin antibiotic resistance ("Amp<sup>r</sup>") and contains a bacterial origin of replication ("ori"), an IPTG inducible promoter, a ribosome binding site ("RBS"), six codons encoding histidine residues that allow affinity purification using nickel-nitrilo-tri-acetic acid ("Ni-NTA") affinity resin sold by QIAGEN, Inc., *supra*, and suitable single restriction enzyme cleavage sites. These elements are arranged such that an inserted DNA fragment encoding a polypeptide expresses that polypeptide with the six His residues (i.e., a "6 X His tag") covalently linked to the carboxyl terminus of that polypeptide.

The DNA sequence encoding the desired portion I-FLICE-1 protein is amplified from the deposited cDNA clone using PCR oligonucleotide primers which anneal to the amino terminal sequences of the desired portion of the I-FLICE-1 protein and to sequences in the deposited construct 3' to the cDNA coding sequence. Additional nucleotides containing restriction sites to facilitate cloning in the pQE60 vector are added to the 5' and 3' sequences, respectively.

For cloning the protein, the 5' primer has the sequence:

5' CGCCCATGGCTGAAGTCATCCATCAG 3' (SEQ ID NO:7) containing the underlined NcoI restriction site followed by 16 (i.e., 275-291) nucleotides complementary to the amino terminal coding sequence of the I-FLICE-1 sequence in FIG. 1A-1B (SEQ ID NO:1). One of ordinary skill in the art would appreciate, of course, that the point in the protein coding sequence where the 5' primer begins may be varied to amplify a DNA segment encoding any desired portion of the complete protein in a shorter or longer form. The 3' primer has the sequence:

5' CGCAAGCTTGTGCTGGGATTACAGGTG 3' (SEQ ID NO:8) containing the underlined HindIII restriction site followed by 18 (i.e., 1740-1758) nucleotides complementary to the 3' end of the coding sequence immediately before the stop codon in the I-FLICE-1 DNA sequence in FIG. 1A-1B (SEQ ID NO:1), with the coding sequence aligned with the restriction site so as to maintain its reading frame with that of the six His codons in the pQE60 vector.

The amplified I-FLICE-1 DNA fragment and the vector pQE60 are digested with NcoI/HindIII and the digested DNAs are then ligated together. Insertion of the I-FLICE-1 DNA into the restricted pQE60 vector places the I-FLICE-1 protein coding region downstream from the IPTG-inducible promoter and in-frame with an initiating AUG and the six histidine codons.

The ligation mixture is transformed into competent *E. coli* cells using standard procedures such as those described in Sambrook *et al.*, *Molecular Cloning: a Laboratory Manual, 2nd Ed.*; Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY (1989). *E. coli* strain M15/rep4, containing multiple copies of the plasmid pREP4, which expresses the lac repressor and confers kanamycin resistance ("Kan<sup>r</sup>"), is used in carrying out the illustrative example described herein. This strain, which is only one of many that are suitable for expressing I-FLICE-1 protein, is available commercially from QIAGEN, Inc., *supra*. Transformants are identified by their ability to grow on LB plates in the presence of ampicillin and kanamycin. Plasmid DNA is isolated from resistant

colonies and the identity of the cloned DNA confirmed by restriction analysis, PCR and DNA sequencing.

Clones containing the desired constructs are grown overnight ("O/N") in liquid culture in LB media supplemented with both ampicillin (100 µg/ml) and kanamycin (25 µg/ml). The O/N culture is used to inoculate a large culture, at a dilution of approximately 1:25 to 1:250. The cells are grown to an optical density at 600 nm ("OD600") of between 0.4 and 0.6. Isopropyl-b-D-thiogalactopyranoside ("IPTG") is then added to a final concentration of 1 mM to induce transcription from the lac repressor sensitive promoter, by inactivating the lacI repressor. Cells subsequently are incubated further for 3 to 4 hours. Cells then are harvested by centrifugation.

The cells are then stirred for 3-4 hours at 4°C in 6M guanidine-HCl, pH 8. The cell debris is removed by centrifugation, and the supernatant containing the I-FLICE-1 is loaded onto a nickel-nitrilo-tri-acetic acid ("NiNTA") affinity resin column (available from QIAGEN, Inc., *supra*). Proteins with a 6 x His tag bind to the Ni-NTA resin with high affinity and can be purified in a simple one-step procedure (for details see: The QIAexpressionist, 1995, QIAGEN, Inc., *supra*). Briefly the supernatant is loaded onto the column in 6 M guanidine-HCl, pH8, the column is first washed with 10 volumes of 6 M guanidine-HCl, pH8, then washed with 10 volumes of 6 M guanidine-HCl pH6, and finally the I-FLICE-1 is eluted with 6 M guanidine-HCl, pH5.

The purified protein is then renatured by dialyzing it against phosphate-buffered saline (PBS) or 50 mM Na-acetate, pH 6 buffer plus 200 mM NaCl. Alternatively, the protein can be successfully refolded while immobilized on the Ni-NTA column. The recommended conditions are as follows: renature using a linear 6M-1M urea gradient in 500 mM NaCl, 20% glycerol, 20 mM Tris/HCl pH7.4, containing protease inhibitors. The renaturation should be performed over a period of 1.5 hours or more. After renaturation the proteins can be eluted by the addition of 250 mM imidazole. Imidazole is removed by a final dialyzing step

-42-

against PBS or 50 mM sodium acetate pH6 buffer plus 200 mM NaCl. The purified protein is stored at 4°C or frozen at -80°C.

***Example 1(b): Expression and Purification of I-FLICE-2 in E. coli***

5 The bacterial expression vector pQE60 is used for bacterial expression in this example. (QIAGEN, Inc., 9259 Eton Avenue, Chatsworth, CA, 91311). pQE60 encodes ampicillin antibiotic resistance ("Amp<sup>r</sup>") and contains a bacterial origin of replication ("ori"), an IPTG inducible promoter, a ribosome binding site ("RBS"), six codons encoding histidine residues that allow affinity purification using nickel-nitrilo-tri-acetic acid ("Ni-NTA") affinity resin sold by QIAGEN,  
10 Inc., *supra*, and suitable single restriction enzyme cleavage sites. These elements are arranged such that an inserted DNA fragment encoding a polypeptide expresses that polypeptide with the six His residues (i.e., a "6 X His tag") covalently linked to the carboxyl terminus of that polypeptide.

15 The DNA sequence encoding the desired portion I-FLICE-2 protein is amplified from the deposited cDNA clone using PCR oligonucleotide primers which anneal to the amino terminal sequences of the desired portion of the I-FLICE-2 protein and to sequences in the deposited construct 3' to the cDNA coding sequence. Additional nucleotides containing restriction sites to facilitate cloning in the pQE60 vector are added to the 5' and 3' sequences, respectively.

20 For cloning the protein, the 5' primer has the sequence:  
5' CGCCCATGGAGATTGGTGAGGATTTG 3' (SEQ ID NO:9) containing the underlined NcoI restriction site followed by 17 (i.e., 311-328) nucleotides complementary to the amino terminal coding sequence of the I-FLICE-2 sequence in FIG. 4A-4C (SEQ ID NO:5). one of ordinary skill in the art would appreciate,  
25 of course, that the point in the protein coding sequence where the 5' primer begins may be varied to amplify a DNA segment encoding any desired portion of the complete protein in a shorter or longer form. The 3' primer has the sequence:



5' CGCAAGCTTAGAGCATGCAGTGTCAG 3' (SEQ ID NO:10) containing the underlined HindIII restriction site followed by 16 (i.e., 1400-1416) nucleotides complementary to the 3' end of the coding sequence immediately before the stop codon in the I-FLICE-2 DNA sequence in FIG. 4A-4C (SEQ ID NO:5), with the coding sequence aligned with the restriction site so as to maintain its reading frame with that of the six His codons in the pQE60 vector.

The amplified I-FLICE-2 DNA fragment and the vector pQE60 are digested with NcoI/HindIII and the digested DNAs are then ligated together. Insertion of the I-FLICE-2 DNA into the restricted pQE60 vector places the I-FLICE-2 protein coding region downstream from the IPTG-inducible promoter and in-frame with an initiating AUG and the six histidine codons.

The ligation mixture is transformed into competent *E. coli* cells using standard procedures such as those described in Sambrook *et al.*, *Molecular Cloning: a Laboratory Manual, 2nd Ed.*; Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY (1989). *E. coli* strain M15/rep4, containing multiple copies of the plasmid pREP4, which expresses the lac repressor and confers kanamycin resistance ("Kan"), is used in carrying out the illustrative example described herein. This strain, which is only one of many that are suitable for expressing I-FLICE-2 protein, is available commercially from QIAGEN, Inc., *supra*. Transformants are identified by their ability to grow on LB plates in the presence of ampicillin and kanamycin. Plasmid DNA is isolated from resistant colonies and the identity of the cloned DNA confirmed by restriction analysis, PCR and DNA sequencing.

Clones containing the desired constructs are grown overnight ("O/N") in liquid culture in LB media supplemented with both ampicillin (100 µg/ml) and kanamycin (25 µg/ml). The O/N culture is used to inoculate a large culture, at a dilution of approximately 1:25 to 1:250. The cells are grown to an optical density at 600 nm ("OD600") of between 0.4 and 0.6. Isopropyl-b-D-thiogalactopyranoside ("IPTG") is then added to a final concentration of 1 mM to induce transcription from the lac repressor sensitive promoter, by inactivating

the lacI repressor. Cells subsequently are incubated further for 3 to 4 hours. Cells then are harvested by centrifugation.

The cells are then stirred for 3-4 hours at 4°C in 6M guanidine-HCl, pH8. The cell debris is removed by centrifugation, and the supernatant containing the I-FLICE-2 is loaded onto a nickel-nitrilo-tri-acetic acid ("NiNTA") affinity resin column (available from QIAGEN, Inc., *supra*). Proteins with a 6 x His tag bind to the Ni-NTA resin with high affinity and can be purified in a simple one-step procedure (for details see: The QIAexpressionist, 1995, QIAGEN, Inc., *supra*). Briefly the supernatant is loaded onto the column in 6 M guanidine-HCl, pH8, the column is first washed with 10 volumes of 6 M guanidine-HCl, pH8, then washed with 10 volumes of 6 M guanidine-HCl pH6, and finally the I-FLICE-2 is eluted with 6 M guanidine-HCl, pH5.

The purified protein is then renatured by dialyzing it against phosphate-buffered saline (PBS) or 50 mM Na-acetate, pH 6 buffer plus 200 mM NaCl. Alternatively, the protein can be successfully refolded while immobilized on the Ni-NTA column. The recommended conditions are as follows: renature using a linear 6M-1M urea gradient in 500 mM NaCl, 20% glycerol, 20 mM Tris/HCl pH7.4, containing protease inhibitors. The renaturation should be performed over a period of 1.5 hours or more. After renaturation the proteins can be eluted by the addition of 250 mM imidazole. Imidazole is removed by a final dialyzing step against PBS or 50 mM sodium acetate pH6 buffer plus 200 mM NaCl. The purified protein is stored at 4°C or frozen at -80°C.

***Example 2(a): Cloning and Expression of I-FLICE-1 protein in a Baculovirus Expression System***

In this illustrative example, the plasmid shuttle vector pA2 is used to insert the cloned DNA encoding the complete protein into a baculovirus to express the I-FLICE-1 protein, using standard methods as described in Summers *et al.*, *A Manual of Methods for Baculovirus Vectors and Insect Cell Culture Procedures*, Texas Agricultural Experimental Station Bulletin No. 1555 (1987).

This expression vector contains the strong polyhedrin promoter of the *Autographa californica* nuclear polyhedrosis virus (AcMNPV) followed by convenient restriction sites such as BamHI and Asp718. The polyadenylation site of the simian virus 40 ("SV40") is used for efficient polyadenylation. For easy selection of recombinant virus, the plasmid contains the beta-galactosidase gene from *E. coli* under control of a weak *Drosophila* promoter in the same orientation, followed by the polyadenylation signal of the polyhedrin gene. The inserted genes are flanked on both sides by viral sequences for cell-mediated homologous recombination with wild-type viral DNA to generate viable virus that express the cloned polynucleotide.

Many other baculovirus vectors could be used in place of the vector above, such as pAc373, pVL941 and pAcIM1, as one skilled in the art would readily appreciate, as long as the construct provides appropriately located signals for transcription, translation, secretion and the like, including a signal peptide and an in-frame AUG as required. Such vectors are described, for instance, in Luckow *et al.*, *Virology* 170:31-39.

The cDNA sequence encoding the full length I-FLICE-1 protein in the deposited clone, including the AUG initiation codon shown in FIG. 1A-1B (SEQ ID NO:1), is amplified using PCR oligonucleotide primers corresponding to the 5' and 3' sequences of the gene. The 5' primer has the sequence:

5' CGCGGATCCGCCATCATGTCTGCTGAAGTCATC 3' (SEQ ID NO:11) containing the underlined BamHI restriction enzyme site, an efficient signal for initiation of translation in eukaryotic cells, as described by Kozak, M., *J. Mol. Biol.* 196:947-950 (1987), followed by 17 (i.e., 268-285) bases of the sequence of the complete I-FLICE-1 protein shown in FIG. 1A-1B, beginning with the AUG initiation codon. The 3' primer has the sequence:

5' CGCGGTACCGTGCTGGGATTACAGGTG 3' (SEQ ID NO:12) containing the underlined, Asp718 restriction site followed by 18 (1740-1758) nucleotides complementary to the 3' noncoding sequence in FIG. 1A-1B (SEQ ID NO:1).

-46-

The amplified fragment is isolated from a 1% agarose gel using a commercially available kit ("GeneClean," BIO 101 Inc., La Jolla, Ca.). The fragment then is digested with BamHI and Asp718 and again is purified on a 1% agarose gel. This fragment is designated herein "F1".

5        The plasmid is digested with the restriction enzymes BamHI and Asp718 and optionally, can be dephosphorylated using calf intestinal phosphatase, using routine procedures known in the art. The DNA is then isolated from a 1% agarose gel using a commercially available kit ("GeneClean" BIO 101 Inc., La Jolla, Ca.). This vector DNA is designated herein "V1".

10        Fragment F1 and the dephosphorylated plasmid V1 are ligated together with T4 DNA ligase. *E. coli* HB101 or other suitable *E. coli* hosts such as XL-1 Blue (Stratagene Cloning Systems, La Jolla, CA) cells are transformed with the ligation mixture and spread on culture plates. Bacteria are identified that contain the plasmid with the human I-FLICE-1 gene using the PCR method, in which one  
15        of the primers that is used to amplify the gene and the second primer is from well within the vector so that only those bacterial colonies containing the I-FLICE-1 gene fragment will show amplification of the DNA. The sequence of the cloned fragment is confirmed by DNA sequencing. This plasmid is designated herein pBac I-FLICE-1.

20        Five  $\mu$ g of the plasmid pBac I-FLICE-1 is co-transfected with 1.0  $\mu$ g of a commercially available linearized baculovirus DNA ("BaculoGold™ baculovirus DNA", Pharmingen, San Diego, CA.), using the lipofection method described by Felgner *et al.*, *Proc. Natl. Acad. Sci. USA* 84:7413-7417 (1987). 1  $\mu$ g of BaculoGold™ virus DNA and 5  $\mu$ g of the plasmid pBac I-FLICE-1 are  
25        mixed in a sterile well of a microtiter plate containing 50  $\mu$ l of serum-free Grace's medium (Life Technologies Inc., Gaithersburg, MD). Afterwards, 10  $\mu$ l Lipofectin plus 90  $\mu$ l Grace's medium are added, mixed and incubated for 15 minutes at room temperature. Then the transfection mixture is added drop-wise to Sf9 insect cells (ATCC CRL 1711) seeded in a 35 mm tissue culture plate with  
30        1 ml Grace's medium without serum. The plate is rocked back and forth to mix

-47-

the newly added solution. The plate is then incubated for 5 hours at 27°C. After 5 hours the transfection solution is removed from the plate and 1 ml of Grace's insect medium supplemented with 10% fetal calf serum is added. The plate is put back into an incubator and cultivation is continued at 27°C for four days.

5           After four days the supernatant is collected and a plaque assay is performed, as described by Summers and Smith, *supra*. An agarose gel with "Blue Gal" (Life Technologies Inc., Gaithersburg) is used to allow easy identification and isolation of gal-expressing clones, which produce blue-stained  
10           plaques. (A detailed description of a "plaque assay" of this type can also be found in the user's guide for insect cell culture and baculovirology distributed by Life Technologies Inc., Gaithersburg, page 9-10). After appropriate incubation, blue stained plaques are picked with the tip of a micropipettor (e.g., Eppendorf). The agar containing the recombinant viruses is then resuspended in a microcentrifuge  
15           tube containing 200 µl of Grace's medium and the suspension containing the recombinant baculovirus is used to infect Sf9 cells seeded in 35 mm dishes. Four days later the supernatants of these culture dishes are harvested and then they are stored at 4°C. The recombinant virus is called V- I-FLICE-1.

          To verify the expression of the I-FLICE-1 gene, Sf9 cells are grown in Grace's medium supplemented with 10% heat inactivated FBS. The cells are  
20           infected with the recombinant baculovirus V- I-FLICE-1 at a multiplicity of infection ("MOI") of about 2. Six hours later the medium is removed and is replaced with SF900 II medium minus methionine and cysteine (available from Life Technologies Inc., Rockville, MD). If radiolabeled proteins are desired, 42  
25           hours later, 5 µCi of <sup>35</sup>S-methionine and 5 µCi <sup>35</sup>S-cysteine (available from Amersham) are added. The cells are further incubated for 16 hours and then they are harvested by centrifugation. The proteins in the supernatant as well as the intracellular proteins are analyzed by SDS-PAGE followed by autoradiography (if radiolabeled). Microsequencing of the amino acid sequence of the amino terminus of purified protein may be used to determine the amino terminal

sequence of the mature protein and thus the cleavage point and length of the secretory signal peptide.

***Example 2(b): Cloning and Expression of I-FLICE-2 protein in a Baculovirus Expression System***

5 In this illustrative example, the plasmid shuttle vector pA2 is used to insert the cloned DNA encoding the complete protein into a baculovirus to express the I-FLICE-2 protein, using standard methods as described in Summers *et al.*, *A Manual of Methods for Baculovirus Vectors and Insect Cell Culture Procedures*, Texas Agricultural Experimental Station Bulletin No. 1555 (1987).  
10 This expression vector contains the strong polyhedrin promoter of the *Autographa californica* nuclear polyhedrosis virus (AcMNPV) followed by convenient restriction sites such as BamHI and Asp718. The polyadenylation site of the simian virus 40 ("SV40") is used for efficient polyadenylation. For easy selection of recombinant virus, the plasmid contains the beta-galactosidase gene  
15 from *E. coli* under control of a weak *Drosophila* promoter in the same orientation, followed by the polyadenylation signal of the polyhedrin gene. The inserted genes are flanked on both sides by viral sequences for cell-mediated homologous recombination with wild-type viral DNA to generate viable virus that express the cloned polynucleotide.

20 Many other baculovirus vectors could be used in place of the vector above, such as pAc373, pVL941 and pAcIM1, as one skilled in the art would readily appreciate, as long as the construct provides appropriately located signals for transcription, translation, secretion and the like, including a signal peptide and an in-frame AUG as required. Such vectors are described, for instance, in  
25 Luckow *et al.*, *Virology* 170:31-39.

The cDNA sequence encoding the full length I-FLICE-2 protein in the deposited clone, including the AUG initiation codon shown in FIG. 4A-4C (SEQ ID NO:6) is amplified using PCR oligonucleotide primers corresponding to the 5' and 3' sequences of the gene. The 5' primer has the sequence:

5' CGCGGATCCGCCATCATGGCAGAGATTGGTGAG 3' (SEQ ID NO:13) containing the underlined BamHI restriction enzyme site, an efficient signal for initiation of translation in eukaryotic cells, as described by Kozak, M., *J. Mol. Biol.* 196:947-950 (1987), followed by 17 (304-321) bases of the sequence of the complete I-FLICE-2 protein shown in FIG. 4A-4C, beginning with the AUG initiation codon. The 3' primer has the sequence:

5' CGCGGTACCAGAGCATGCAGTGTCTCAG 3' (SEQ ID NO:14) containing the underlined, Asp718 restriction site followed by (i.e., 1400-1416) nucleotides complementary to the 3' noncoding sequence in FIG. 4A-4C (SEQ ID NO:5).

The amplified fragment is isolated from a 1% agarose gel using a commercially available kit ("GeneClean," BIO 101 Inc., La Jolla, Ca.). The fragment then is digested with BamHI and Asp718 and again is purified on a 1% agarose gel. This fragment is designated herein "F1".

The plasmid is digested with the restriction enzymes BamHI and Asp718 and optionally, can be dephosphorylated using calf intestinal phosphatase, using routine procedures known in the art. The DNA is then isolated from a 1% agarose gel using a commercially available kit ("GeneClean" BIO 101 Inc., La Jolla, Ca.). This vector DNA is designated herein "V1".

Fragment F1 and the dephosphorylated plasmid V1 are ligated together with T4 DNA ligase. *E. coli* HB101 or other suitable *E. coli* hosts such as XL-1 Blue (Stratagene Cloning Systems, La Jolla, CA) cells are transformed with the ligation mixture and spread on culture plates. Bacteria are identified that contain the plasmid with the human I-FLICE-2 gene using the PCR method, in which one of the primers that is used to amplify the gene and the second primer is from well within the vector so that only those bacterial colonies containing the I-FLICE-2 gene fragment will show amplification of the DNA. The sequence of the cloned fragment is confirmed by DNA sequencing. This plasmid is designated herein pBac I-FLICE-2.

Five  $\mu$ g of the plasmid pBac I-FLICE-2 is co-transfected with 1.0  $\mu$ g of a commercially available linearized baculovirus DNA ("BaculoGold™

5 baculovirus DNA", Pharmingen, San Diego, CA.), using the lipofection method described by Felgner *et al.*, *Proc. Natl. Acad. Sci. USA* 84:7413-7417 (1987). 1 µg of BaculoGold™ virus DNA and 5 µg of the plasmid pBac I-FLICE-2 are mixed in a sterile well of a microtiter plate containing 50 µl of serum-free Grace's medium (Life Technologies Inc., Gaithersburg, MD). Afterwards, 10 µl Lipofectin plus 90 µl Grace's medium are added, mixed and incubated for 15 minutes at room temperature. Then the transfection mixture is added drop-wise to Sf9 insect cells (ATCC CRL 1711) seeded in a 35 mm tissue culture plate with 1 ml Grace's medium without serum. The plate is rocked back and forth to mix the newly added solution. The plate is then incubated for 5 hours at 27°C. After 10 5 hours the transfection solution is removed from the plate and 1 ml of Grace's insect medium supplemented with 10% fetal calf serum is added. The plate is put back into an incubator and cultivation is continued at 27°C for four days.

15 After four days the supernatant is collected and a plaque assay is performed, as described by Summers and Smith, *supra*. An agarose gel with "Blue Gal" (Life Technologies Inc., Gaithersburg) is used to allow easy identification and isolation of gal-expressing clones, which produce blue-stained plaques. (A detailed description of a "plaque assay" of this type can also be found in the user's guide for insect cell culture and baculovirology distributed by Life Technologies Inc., Gaithersburg, page 9-10). After appropriate incubation, blue stained plaques are picked with the tip of a micropipettor (e.g., Eppendorf). The agar containing the recombinant viruses is then resuspended in a microcentrifuge tube containing 200 µl of Grace's medium and the suspension containing the recombinant baculovirus is used to infect Sf9 cells seeded in 35 mm dishes. Four 20 days later the supernatants of these culture dishes are harvested and then they are stored at 4°C. The recombinant virus is called V-I-FLICE-2.

25 To verify the expression of the I-FLICE-2 gene, Sf9 cells are grown in Grace's medium supplemented with 10% heat inactivated FBS. The cells are infected with the recombinant baculovirus V-I-FLICE-2 at a multiplicity of infection ("MOI") of about 2. Six hours later the medium is removed and is 30



-51-

replaced with SF900 II medium minus methionine and cysteine (available from Life Technologies Inc., Rockville, MD). If radiolabeled proteins are desired, 42 hours later, 5  $\mu$ Ci of  $^{35}$ S-methionine and 5  $\mu$ Ci  $^{35}$ S-cysteine (available from Amersham) are added. The cells are further incubated for 16 hours and then they are harvested by centrifugation. The proteins in the supernatant as well as the intracellular proteins are analyzed by SDS-PAGE followed by autoradiography (if radiolabeled). Microsequencing of the amino acid sequence of the amino terminus of purified protein may be used to determine the amino terminal sequence of the mature protein and thus the cleavage point and length of the secretory signal peptide.

***Example 3: Cloning and Expression of I-FLICE in Mammalian Cells***

A typical mammalian expression vector contains the promoter element, which mediates the initiation of transcription of mRNA, the protein coding sequence, and signals required for the termination of transcription and polyadenylation of the transcript. Additional elements include enhancers, Kozak sequences and intervening sequences flanked by donor and acceptor sites for RNA splicing. Highly efficient transcription can be achieved with the early and late promoters from SV40, the long terminal repeats (LTRS) from Retroviruses, e.g., RSV, HTLV, HIV and the early promoter of the cytomegalovirus (CMV). However, cellular elements can also be used (e.g., the human actin promoter). Suitable expression vectors for use in practicing the present invention include, for example, vectors such as PSVL and PMSG (Pharmacia, Uppsala, Sweden), pRSVcat (ATCC 37152), pSV2dhfr (ATCC 37146) and pBC12MI (ATCC 67109). Mammalian host cells that could be used include, human HeLa 293, H9 and Jurkat cells, mouse NIH3T3 and C127 cells, Cos 1, Cos 7 and CV 1, quail QC1-3 cells, mouse L cells and Chinese hamster ovary (CHO) cells.

Alternatively, the gene can be expressed in stable cell lines that contain the gene integrated into a chromosome. The co-transfection with a selectable

marker such as dhfr, gpt, neomycin, or hygromycin allows the identification and isolation of the transfected cells.

The transfected gene can also be amplified to express large amounts of the encoded protein. The DHFR (dihydrofolate reductase) marker is useful to develop cell lines that carry several hundred or even several thousand copies of the gene of interest. Another useful selection marker is the enzyme glutamine synthase (GS) (Murphy *et al.*, *Biochem J.* 227:277-279 (1991); Bebbington *et al.*, *Bio/Technology* 10:169-175 (1992)). Using these markers, the mammalian cells are grown in selective medium and the cells with the highest resistance are selected. These cell lines contain the amplified gene(s) integrated into a chromosome. Chinese hamster ovary (CHO) and NSO cells are often used for the production of proteins.

The expression vectors pC1 and pC4 contain the strong promoter (LTR) of the Rous Sarcoma Virus (Cullen *et al.*, *Molec. Cell. Biol.* 5:438-447 (1985)) plus a fragment of the CMV-enhancer (Boshart *et al.*, *Cell* 41:521-530 (1985)). Multiple cloning sites, e.g., with the restriction enzyme cleavage sites BamHI, XbaI and Asp718, facilitate the cloning of the gene of interest. The vectors contain in addition the 3' intron, the polyadenylation and termination signal of the rat preproinsulin gene.

### ***Example 3(a): Cloning and Expression of I-FLICE-1 in COS Cells***

The expression plasmid, p I-FLICE-1 HA, is made by cloning a cDNA encoding I-FLICE-1 into the expression vector pcDNAI/Amp or pcDNAIII (which can be obtained from Invitrogen, Inc.).

The expression vector pcDNAI/amp contains: (1) an *E. coli* origin of replication effective for propagation in *E. coli* and other prokaryotic cells; (2) an ampicillin resistance gene for selection of plasmid-containing prokaryotic cells; (3) an SV40 origin of replication for propagation in eukaryotic cells; (4) a CMV promoter, a polylinker, an SV40 intron; (5) several codons encoding a

hemagglutinin fragment (i.e., an "HA" tag to facilitate purification) followed by a termination codon and polyadenylation signal arranged so that a cDNA can be conveniently placed under expression control of the CMV promoter and operably linked to the SV40 intron and the polyadenylation signal by means of restriction sites in the polylinker. The HA tag corresponds to an epitope derived from the influenza hemagglutinin protein described by Wilson *et al.*, *Cell* 37:767-778 (1984). The fusion of the HA tag to the target protein allows easy detection and recovery of the recombinant protein with an antibody that recognizes the HA epitope. pcDNAIII contains, in addition, the selectable neomycin marker.

A DNA fragment encoding the I-FLICE-1 is cloned into the polylinker region of the vector so that recombinant protein expression is directed by the CMV promoter. The plasmid construction strategy is as follows. The I-FLICE-1 cDNA of the deposited clone is amplified using primers that contain convenient restriction sites, much as described above for construction of vectors for expression of I-FLICE-1 in *E. coli*. Suitable primers include the following, which are used in this example. The 5' primer, containing the underlined SmaI site, a Kozak sequence, an AUG start codon and 17 bases of the 5' coding region of the complete I-FLICE-1 has the following sequence:

5' CGCCCCGGGGCCATCATGTCTGCTGAAGTCATC (268-285) 3' (SEQ ID NO:15). The 3' primer, containing the underlined XbaI site, a stop codon, and 18 bp of 3' coding sequence has the following sequence (at the 3' end):

5' CGCTCTAGATCAAGCGTAGTCTGGGACGTCGTATGGGTAGTGC TGGGATTACAGGTG (1740-1758) 3' (SEQ ID NO:16).

The PCR amplified DNA fragment and the vector, pcDNAI/Amp, are digested with SmaI and XbaI and then ligated. The ligation mixture is transformed into *E. coli* strain SURE (available from Stratagene Cloning Systems, 11099 North Torrey Pines Road, La Jolla, CA 92037), and the -transformed culture is plated on ampicillin media plates which then are incubated to allow growth of ampicillin resistant colonies. Plasmid DNA is isolated from

resistant colonies and examined by restriction analysis or other means for the presence of the I-FLICE-1-encoding fragment.

For expression of recombinant I-FLICE-1, COS cells are transfected with an expression vector, as described above, using DEAE-DEXTRAN, as described, for instance, in Sambrook *et al.*, *Molecular Cloning: a Laboratory Manual*, Cold Spring Laboratory Press, Cold Spring Harbor, New York (1989). Cells are incubated under conditions for expression of I-FLICE-1 by the vector.

Expression of the I-FLICE-1-HA fusion protein is detected by radiolabeling and immunoprecipitation, using methods described in, for example Harlow *et al.*, *Antibodies: A Laboratory Manual, 2nd Ed.*; Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (1988). To this end, two days after transfection, the cells are labeled by incubation in media containing <sup>35</sup>S-cysteine for 8 hours. The cells and the media are collected, and the cells are washed and lysed with detergent-containing RIPA buffer: 150 mM NaCl, 1% NP-40, 0.1% SDS, 0.5% DOC, 50 mM TRIS, pH 7.5, as described by Wilson *et al.* cited above. Proteins are precipitated from the cell lysate and from the culture media using an HA-specific monoclonal antibody. The precipitated proteins then are analyzed by SDS-PAGE and autoradiography. An expression product of the expected size is seen in the cell lysate, which is not seen in negative controls.

### ***Example 3(b): Cloning and Expression of I-FLICE-1 in CHO Cells***

The vector pC4 is used for the expression of I-FLICE-1 protein. Plasmid pC4 is a derivative of the plasmid pSV2-dhfr (ATCC Accession No. 37146). The plasmid contains the mouse DHFR gene under control of the SV40 early promoter. Chinese hamster ovary- or other cells lacking dihydrofolate activity that are transfected with these plasmids can be selected by growing the cells in a selective medium (alpha minus MEM, Life Technologies) supplemented with the chemotherapeutic agent methotrexate. The amplification of the DHFR genes in cells resistant to methotrexate (MTX) has been well documented (see, e.g., Alt,

F. W., Kellems, R. M., Bertino, J. R., and Schimke, R. T., 1978, *J Biol. Chem.* 253:1357-1370, Hamlin, J. L. and Ma, C. 1990, *Biochem. et Biophys. Acta*, 1097:107-143, Page, M. J. and Sydenham, M.A. 1991, *Biotechnology* 9:64-68).

Cells grown in increasing concentrations of MTX develop resistance to the drug by overproducing the target enzyme, DHFR, as a result of amplification of the DHFR gene. If a second gene is linked to the DHFR gene, it is usually co-amplified and over-expressed. It is known in the art that this approach may be used to develop cell lines carrying more than 1,000 copies of the amplified gene(s). Subsequently, when the methotrexate is withdrawn, cell lines are obtained which contain the amplified gene integrated into one or more chromosome(s) of the host cell.

Plasmid pC4 contains for expressing the gene of interest the strong promoter of the long terminal repeat (LTR) of the Rous Sarcoma Virus (Cullen *et al.*, *Molec. Cell. Biol.* 5:438-447 (1985)) plus a fragment isolated from the enhancer of the immediate early gene of human cytomegalovirus (CMV) (Boshart *et al.*, *Cell* 41:521-530 (1985)). Downstream of the promoter are BamHI, XbaI, and Asp718 restriction enzyme cleavage sites that allow integration of the genes. Behind these cloning sites the plasmid contains the 3' intron and polyadenylation site of the rat preproinsulin gene. Other high efficiency promoters can also be used for the expression, e.g., the human  $\beta$ -actin promoter, the SV40 early or late promoters or the long terminal repeats from other retroviruses, e.g., HIV and HTLV. Clontech's Tet-Off and Tet-On gene expression systems and similar systems can be used to express the I-FLICE-1 in a regulated way in mammalian cells (Gossen, M., & Bujard, H. 1992, *Proc. Natl. Acad. Sci. USA* 89: 5547-5551). For the polyadenylation of the mRNA other signals, e.g., from the human growth hormone or globin genes can be used as well. Stable cell lines carrying a gene of interest integrated into the chromosomes can also be selected upon co-transfection with a selectable marker such as gpt, G418 or hygromycin. It is advantageous to use more than one selectable marker in the beginning, e.g., G418 plus methotrexate.

The plasmid pC4 is digested with the restriction enzymes BamHI and Asp718 and then dephosphorylated using calf intestinal phosphatase by procedures known in the art. The vector is then isolated from a 1% agarose gel.

The DNA sequence encoding the complete I-FLICE-1 protein including its leader sequence is amplified using PCR oligonucleotide primers corresponding to the 5' and 3' sequences of the gene. The 5' primer has the sequence:

5' CGCGGATCCGCCATCATGTCTGCTGAAGTCATC 3' (SEQ ID NO:17) containing the underlined BamHI restriction enzyme site, an efficient signal for initiation of translation in eukaryotic cells, as described by Kozak, M., *J. Mol. Biol.* 196:947-950 (1987), followed by 17 (i.e., 268-285) bases of the sequence of the complete I-FLICE-1 protein shown in FIG. 1A-1B, beginning with the AUG initiation codon. The 3' primer has the sequence:

5' CGCGGTACCGTGCTGGGATTACAGGTG 3' (SEQ ID NO:18) containing the underlined, Asp718 restriction site followed by 18 (1740-1758) nucleotides complementary to the 3' noncoding sequence in FIG. 1A-1B (SEQ ID NO:1).

The amplified fragment is digested with the endonucleases BamHI and Asp718 and then purified again on a 1% agarose gel. The isolated fragment and the dephosphorylated vector are then ligated with T4 DNA ligase. *E. coli* HB101 or XL-1 Blue cells are then transformed and bacteria are identified that contain the fragment inserted into plasmid pC4 using, for instance, restriction enzyme analysis.

Chinese hamster ovary cells lacking an active DHFR gene are used for transfection. Five µg of the expression plasmid pC4 is cotransfected with 0.5 µg of the plasmid pSV2-neo using lipofectin (Felgner *et al.*, *supra*). The plasmid pSV2neo contains a dominant selectable marker, the neo gene from Tn5 encoding an enzyme that confers resistance to a group of antibiotics including G418. The cells are seeded in alpha minus MEM supplemented with 1 mg/ml G418. After 2 days, the cells are trypsinized and seeded in hybridoma cloning plates (Greiner, Germany) in alpha minus MEM supplemented with 10, 25, or 50 ng/ml of methotrexate plus 1 mg/ml G418. After about 10-14 days single clones are

trypsinized and then seeded in 6-well petri dishes or 10 ml flasks using different concentrations of methotrexate (50 nM, 100 nM, 200 nM, 400 nM, 800 nM). Clones growing at the highest concentrations of methotrexate are then transferred to new 6-well plates containing even higher concentrations of methotrexate (1  $\mu$ M, 2  $\mu$ M, 5  $\mu$ M, 10  $\mu$ M, 20  $\mu$ M). The same procedure is repeated until clones are obtained which grow at a concentration of 100 - 200  $\mu$ M. Expression of the desired gene product is analyzed, for instance, by SDS-PAGE and Western blot or by reverse phase HPLC analysis.

***Example 3(c): Cloning and Expression of I-FLICE-2 in COS Cells***

The expression plasmid, pI-FLICE-2HA, is made by cloning a cDNA encoding I-FLICE-2 into the expression vector pcDNAI/Amp or pcDNAIII (which can be obtained from Invitrogen, Inc.).

The expression vector pcDNAI/amp contains: (1) an *E. coli* origin of replication effective for propagation in *E. coli* and other prokaryotic cells; (2) an ampicillin resistance gene for selection of plasmid-containing prokaryotic cells; (3) an SV40 origin of replication for propagation in eukaryotic cells; (4) a CMV promoter, a polylinker, an SV40 intron; (5) several codons encoding a hemagglutinin fragment (i.e., an "HA" tag to facilitate purification) followed by a termination codon and polyadenylation signal arranged so that a cDNA can be conveniently placed under expression control of the CMV promoter and operably linked to the SV40 intron and the polyadenylation signal by means of restriction sites in the polylinker. The HA tag corresponds to an epitope derived from the influenza hemagglutinin protein described by Wilson *et al.*, *Cell* 37:767-778 (1984). The fusion of the HA tag to the target protein allows easy detection and recovery of the recombinant protein with an antibody that recognizes the HA epitope. pcDNAIII contains, in addition, the selectable neomycin marker.

A DNA fragment encoding the I-FLICE-2 is cloned into the polylinker region of the vector so that recombinant protein expression is directed by the

CMV promoter. The plasmid construction strategy is as follows. The I-FLICE-2 cDNA of the deposited clone is amplified using primers that contain convenient restriction sites, much as described above for construction of vectors for expression of I-FLICE-2 in *E. coli*. Suitable primers include the following, which are used in this example. The 5' primer, containing the underlined BamHI site, a Kozak sequence, an AUG start codon and 17 codons of the 5' coding region of the complete I-FLICE-2 has the following sequence:

5' CGCGGATCCGCCATCATGGCAGAGATTGGTGAG 3' (SEQ ID NO:19).

The 3' primer, containing the underlined XbaI site, a stop codon, and 16 bp of 3' coding sequence has the following sequence (at the 3' end):

5' CGCTCTAGATCAAGCGTAGTCTGGGACGTCGTATGGGTAAGA  
GCATGCAGTGTCAG 3' (SEQ ID NO:20).

The PCR amplified DNA fragment and the vector, pcDNAI/Amp, are digested with BamHI and XbaI and then ligated. The ligation mixture is transformed into *E. coli* strain SURE (available from Stratagene Cloning Systems, 11099 North Torrey Pines Road, La Jolla, CA 92037), and the -transformed culture is plated on ampicillin media plates which then are incubated to allow growth of ampicillin resistant colonies. Plasmid DNA is isolated from resistant colonies and examined by restriction analysis or other means for the presence of the I-FLICE-2-encoding fragment.

For expression of recombinant I-FLICE-2, COS cells are transfected with an expression vector, as described above, using DEAE-DEXTRAN, as described, for instance, in Sambrook *et al.*, *Molecular Cloning: a Laboratory Manual*, Cold Spring Laboratory Press, Cold Spring Harbor, New York (1989). Cells are incubated under conditions for expression of I-FLICE-2 by the vector.

Expression of the I-FLICE-2-HA fusion protein is detected by radiolabeling and immunoprecipitation, using methods described in, for example Harlow *et al.*, *Antibodies: A Laboratory Manual, 2nd Ed.*; Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (1988). To this end, two days after transfection, the cells are labeled by incubation in media containing <sup>35</sup>S-



cysteine for 8 hours. The cells and the media are collected, and the cells are washed and lysed with detergent-containing RIPA buffer: 150 mM NaCl, 1% NP-40, 0.1% SDS, 0.5% DOC, 50 mM TRIS, pH 7.5, as described by Wilson *et al.* cited above. Proteins are precipitated from the cell lysate and from the culture media using an HA-specific monoclonal antibody. The precipitated proteins then are analyzed by SDS-PAGE and autoradiography. An expression product of the expected size is seen in the cell lysate, which is not seen in negative controls.

***Example 3(d): Cloning and Expression of I-FLICE-2 in CHO Cells***

The vector pC4 is used for the expression of I-FLICE-2 protein. Plasmid pC4 is a derivative of the plasmid pSV2-dhfr (ATCC Accession No. 37146). The plasmid contains the mouse DHFR gene under control of the SV40 early promoter. Chinese hamster ovary- or other cells lacking dihydrofolate activity that are transfected with these plasmids can be selected by growing the cells in a selective medium (alpha minus MEM, Life Technologies) supplemented with the chemotherapeutic agent methotrexate. The amplification of the DHFR genes in cells resistant to methotrexate (MTX) has been well documented (see, e.g., Alt, F. W., Kellems, R. M., Bertino, J. R., and Schimke, R. T., 1978, *J Biol. Chem.* 253:1357-1370, Hamlin, J. L. and Ma, C. 1990, *Biochem. et Biophys. Acta*, 1097:107-143, Page, M. J. and Sydenham, M.A. 1991, *Biotechnology* 9:64-68). Cells grown in increasing concentrations of MTX develop resistance to the drug by overproducing the target enzyme, DHFR, as a result of amplification of the DHFR gene. If a second gene is linked to the DHFR gene, it is usually co-amplified and over-expressed. It is known in the art that this approach may be used to develop cell lines carrying more than 1,000 copies of the amplified gene(s). Subsequently, when the methotrexate is withdrawn, cell lines are obtained which contain the amplified gene integrated into one or more chromosome(s) of the host cell.

Plasmid pC4 contains for expressing the gene of interest the strong promoter of the long terminal repeat (LTR) of the Rous Sarcoma Virus (Cullen *et al.*, *Molec. Cell. Biol.* 5:438-447 (1985)) plus a fragment isolated from the enhancer of the immediate early gene of human cytomegalovirus (CMV) (Boshart *et al.*, *Cell* 41:521-530 (1985)). Downstream of the promoter are BamHI, XbaI, and Asp718 restriction enzyme cleavage sites that allow integration of the genes. Behind these cloning sites the plasmid contains the 3' intron and polyadenylation site of the rat preproinsulin gene. Other high efficiency promoters can also be used for the expression, e.g., the human  $\beta$ -actin promoter, the SV40 early or late promoters or the long terminal repeats from other retroviruses, e.g., HIV and HTLV. Clontech's Tet-Off and Tet-On gene expression systems and similar systems can be used to express the I-FLICE-2 in a regulated way in mammalian cells (Gossen, M., & Bujard, H. 1992, *Proc. Natl. Acad. Sci. USA* 89:5547-5551). For the polyadenylation of the mRNA other signals, e.g., from the human growth hormone or globin genes can be used as well. Stable cell lines carrying a gene of interest integrated into the chromosomes can also be selected upon co-transfection with a selectable marker such as gpt, G418 or hygromycin. It is advantageous to use more than one selectable marker in the beginning, e.g., G418 plus methotrexate.

The plasmid pC4 is digested with the restriction enzymes BamHI/Asp718 and then dephosphorylated using calf intestinal phosphatase by procedures known in the art. The vector is then isolated from a 1% agarose gel.

The DNA sequence encoding the complete I-FLICE-2 protein sequence is amplified using PCR oligonucleotide primers corresponding to the 5' and 3' sequences of the gene. The 5' primer has the sequence:

5' CGCGGATCCGCCATCATGGCAGAGATTGGTGAG 3' (SEQ ID NO:21) containing the underlined BamHI restriction enzyme site, an efficient signal for initiation of translation in eukaryotic cells, as described by Kozak, M., *J. Mol. Biol.* 196:947-950 (1987), followed by 17 (304-321) bases of the sequence of the

complete I-FLICE-2 protein shown in FIG. 4A-4C, beginning with the AUG initiation codon. The 3' primer has the sequence:

5' CGCGGTACCAGAGCATGCAGTGTCTAG 3' (SEQ ID NO:22) containing the underlined, Asp718 restriction site followed by (i.e., 1400-1416) nucleotides complementary to the 3' noncoding sequence in FIG. 4A-4C (SEQ ID NO:5).

The amplified fragment is digested with the endonucleases BamHI and Asp718 and then purified again on a 1% agarose gel. The isolated fragment and the dephosphorylated vector are then ligated with T4 DNA ligase. *E. coli* HB101 or XL-1 Blue cells are then transformed and bacteria are identified that contain the fragment inserted into plasmid pC4 using, for instance, restriction enzyme analysis.

Chinese hamster ovary cells lacking an active DHFR gene are used for transfection. Five  $\mu$ g of the expression plasmid pC4 is cotransfected with 0.5  $\mu$ g of the plasmid pSV2-neo using lipofectin (Felgner *et al.*, *supra*). The plasmid pSV2neo contains a dominant selectable marker, the neo gene from Tn5 encoding an enzyme that confers resistance to a group of antibiotics including G418. The cells are seeded in alpha minus MEM supplemented with 1 mg/ml G418. After 2 days, the cells are trypsinized and seeded in hybridoma cloning plates (Greiner, Germany) in alpha minus MEM supplemented with 10, 25, or 50 ng/ml of methotrexate plus 1 mg/ml G418. After about 10-14 days single clones are trypsinized and then seeded in 6-well petri dishes or 10 ml flasks using different concentrations of methotrexate (50 nM, 100 nM, 200 nM, 400 nM, 800 nM). Clones growing at the highest concentrations of methotrexate are then transferred to new 6-well plates containing even higher concentrations of methotrexate (1  $\mu$ M, 2  $\mu$ M, 5  $\mu$ M, 10  $\mu$ M, 20  $\mu$ M). The same procedure is repeated until clones are obtained which grow at a concentration of 100 - 200  $\mu$ M. Expression of the desired gene product is analyzed, for instance, by SDS-PAGE and Western blot or by reverse phase HPLC analysis.

**Example 4(a): Tissue distribution of I-FLICE-1 mRNA expression**

Northern blot analysis was carried out to examine I-FLICE-1 gene expression in human tissues, using methods described by, among others, Sambrook *et al.*, cited above. A cDNA probe containing the entire nucleotide sequence of the I-FLICE-1 protein (SEQ ID NO:1) was labeled with  $^{32}\text{P}$  using the rediprime™ DNA labeling system (Amersham Life Science), according to manufacturer's instructions. After labeling, the probe was purified using a CHROMA SPIN- 100™ column (Clontech Laboratories, Inc.), according to manufacturer's protocol number PT1200-1. The purified labeled probe was then used to examine various human tissues for I-FLICE-1 mRNA.

Multiple Tissue Northern (MTN) blots containing various human tissues (H) or human immune system tissues (IM) were obtained from Clontech and were examined with the labeled probe using ExpressHyb™ hybridization solution (Clontech) according to manufacturer's protocol number PT1190-1. Following hybridization and washing, the blots were mounted and exposed to film at  $-70^{\circ}\text{C}$  overnight, and films developed according to standard procedures.

Two transcripts were observed (7.5 kb and 6 kb) which presumably represent mRNA sequences encoding I-FLICE-1 and I-FLICE-2. I-FLICE expression was identified in most tissues and cell lines examined except for the brain and the lymphoblastic leukemia line MOLT4. In particular, I-FLICE expression was evident in peripheral blood leukocytes, spleen, placenta and heart.

**Example 4(b): Tissue distribution of I-FLICE-2 mRNA expression**

Northern blot analysis is carried out to examine I-FLICE-2 gene expression in human tissues, using methods described by, among others, Sambrook *et al.*, cited above. A cDNA probe containing the entire nucleotide sequence of the I-FLICE-2 protein (SEQ ID NO:6) is labeled with  $^{32}\text{P}$  using the rediprime™ DNA labeling system (Amersham Life Science), according to

manufacturer's instructions. After labeling, the probe is purified using a CHROMA SPIN- 100™ column (Clontech Laboratories, Inc.), according to manufacturer's protocol number PT1200-1. The purified labeled probe is then used to examine various human tissues for I-FLICE-2 mRNA.

5 Multiple Tissue Northern (MTN) blots containing various human tissues (H) or human immune system tissues (IM) are obtained from Clontech and are examined with the labeled probe using ExpressHyb™ hybridization solution (Clontech) according to manufacturer's protocol number PT1190-1. Following hybridization and washing, the blots are mounted and exposed to film at -70°C  
10 overnight, and films developed according to standard procedures.

***Example 5: I-FLICE-1 Associates with FLICE and Mch4/FLICE-2***

Previous studies have shown that the DED domain is a protein interaction motif that mediates the binding of the adaptor molecule FADD to the effector proteases FLICE and Mch4/FLICE2 (Muzio *et al.*, *Cell* 85:817-27 (1996);  
15 Chinnaiyan *et al.*, *Cell* 81:505-12 (1995)). Given the striking structural similarity, the following experiment was performed to determine whether I-FLICE-1 interacted with either FADD or other FLICE-like caspases.

***Materials and Methods***

*Cell Lines and Expression Vectors* - Human embryonic kidney 293, 293T  
20 and 293-EBNA cells were cultured to Dulbecco's modified Eagle's medium containing 10% fetal bovine serum, nonessential amino acids, L-glutamine, and penicillin/streptomycin. Expression constructs were made in pcDNA3 or pcDNA3.1/MycHisA (Invitrogen) using standard recombinant methodologies (Sambrook, J. *et al.*, *Molecular Cloning* 2nd Edition, Cold Spring Harbor  
25 Laboratory Press).

*Cloning of I-FLICE-1* - cDNAs corresponding to the partial open reading frame of I-FLICE-1 were identified as sequences homologous to FLICE and Mch4/FLICE2 on searching the Human Genome Sciences data base using established EST methods (Adams, M.D. *et al.*, *Science* 252:1651-1656 (1991) and Adams, M.D. *et al.*, *Nature* 355:632-634 (1992)). Full length cDNAs were obtained by screening a random-primed human umbilical vein endothelial cell cDNA library constructed in the pcDNA1 vector (Invitrogen). The sequence of I-FLICE-1 was confirmed by sequencing plasmid DNA template on both strands by the dideoxy chain termination method employing modified T7 DNA polymerase (Sequenase, U.S. Biochemical Corp.).

*Transfection, Coimmunoprecipitation and Western Analysis* - Transient transfections of 293T cells were performed as described previously (O'Rourke *et al.*, *J. Biol. Chem.* 267:24921-24924 (1992)). Cells were harvested 40 hour following transfection, immunoprecipitation with  $\alpha$ -FLAG or  $\alpha$ myc antibodies and analyzed by immunoblotting.

### ***Results and Discussion***

Sequence analysis of a full length cDNA revealed a 1443-base pair open reading frame that encoded a novel protein with a predicated molecular mass of 55.3 kDa (FIG. 1A-1B). Given that the protein had striking homology to FLICE and Mch4/FLICE2 but lacked an active site, making it a potential dominant negative inhibitor, it was designated I-FLICE (for inhibitor of FLICE).

The architecture of I-FLICE-1 was strikingly similar to that of FLICE and Mch4/FLICE2, including two N-terminal DED-like tandem repeats and a region that resembled the caspase catalytic domain. Importantly, I-FLICE-1 did not contain the catalytic cysteine that is normally embedded in the conserved pentapeptide QACRG or QACQG motif present in all known caspases. Rather, the pentapeptide sequence was QNYVV. In addition, based on the x-ray crystal structure of caspase-1 (and caspase-3), amino acid residues His<sup>237</sup> (His<sup>121</sup>), Gly<sup>238</sup>

(Gly<sup>122</sup>), and Cys<sup>285</sup> (Cys<sup>163</sup>) are involved in catalysis, while residues Arg<sup>179</sup> (Arg<sup>64</sup>), Gln<sup>283</sup> (Gln<sup>161</sup>), Arg<sup>341</sup> (Arg<sup>207</sup>), and Ser<sup>347</sup> (Ser<sup>213</sup>) form a binding pocket for the carboxylate side chain of the P1 aspartic acid (Wilson, K.P. *et al.*, *Nature* 370:270-274 (1994), Rotonda, J. *et al.*, *Nat. Struct. Biol.* 3:619-625 (1996), and Fraser, A. *et al.*, *Cell* 85:781-784 (1996)). These seven residues are conserved in all caspases, but only three of them (Gly, Gln, and Ser) are found in I-FLICE-1. Given this lack of conservation of key residues involved in catalysis and substrate binding it can be concluded that I-FLICE-1 is not a cysteine protease and is incapable of binding Asp at the P1 position. Interestingly, the DED domain of I-FLICE-1 was more related to the corresponding domains present in the viral DED-containing inhibitors K13, MC159, and E8, sharing 34%, 31%, and 33% identity (56%, 51%, and 44% similarity), respectively (Hu, S. *et al.*, *J. Biol. Chem.* 272:9621-9624 (1997) and Thorne, M. *et al.*, *Nature* 386:517-521 (1997)).

Co-immunoprecipitation analysis revealed the ability of I-FLICE-1 to bind FLICE and Mch4/FLICE2 but not FADD. In this respect, I-FLICE-1 resembles the viral DED-containing molecule E8 in that it binds FLICE but not FADD (Hu *et al.*, *J. Biol. Chem.* 272:9621-9624 (1997); Bertin *et al.*, *Proc. Natl. Acad. Sci.* 94:1172-1176 (1997)). Since there was no association between I-FLICE-1 and FADD, I-FLICE-1 was not recruited to the CD-95 or TNFR-1 signaling complex as evidenced by its inability to co-precipitate with these receptors.

#### ***Example 6: Cell Death Assay***

Given the ability of the catalytically inactive I-FLICE-1 to complex with FLICE-like caspases, the inventors reasoned that I-FLICE-1 may be acting as a dominant negative inhibitor since the active form of all caspases is a tetramer derived from the processing of two zymogen forms to a four-chain assembly. It follows that a catalytically inert zymogen, such as I-FLICE-1, would be processed to inactive subunits that would result in the generation of a nonfunctional tetrameric protease. This mechanism predicts that I-FLICE-1 should inhibit

TNFR-1 and CD-95-induced apoptosis where FLICE-like caspases play an initiating role. The following cell death assay was performed.

### ***Materials and Methods***

*Cell Death Assay* - Human embryonic kidney 293 (for TNFR-1 killing) or 293 EBNA cells (for CD-95 killing) were transiently transfected with 0.1  $\mu$ g of the reporter plasmid pCMV  $\beta$ -galactosidase plus 0.5  $\mu$ g of test plasmid in the presence or absence of 2.0  $\mu$ g of inhibitory plasmids. 22-24 hours after transfection, cells were fixed in 0.5% glutaraldehyde and stained with X-gal. Percentage of apoptotic cells was determined by calculating the fraction of membrane blebbed blue cells as a function of total blue cells. All assays were evaluated in duplicate and the mean and the standard deviation calculated.

### ***Results***

Consistent with the proposed mechanism, overexpression of I-FLICE-1 resulted in substantial inhibition of TNFR-1 induced cell death comparable to previously characterized inhibitors including CrmA, MC159, dominant negative FLICE (DNFLICE) and Mch4/FLICE2 (DNFLICE2) (see FIG. 6A). However, under the present experimental conditions, I-FLICE-1 appeared to be a less potent inhibitor of CD-95 induced cell death, possibly reflecting the more potent death signal that emanates from this receptor (see FIG. 6B).

It will be clear that the invention may be practiced otherwise than as particularly described in the foregoing description and examples. Numerous modifications and variations of the present invention are possible in light of the above teachings and, therefore, are within the scope of the appended claims.

The entire disclosure of all publications (including patents, patent applications, journal articles, laboratory manuals, books, or other documents) cited herein are hereby incorporated by reference.



- 67 -

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

- (i) APPLICANTS: HUMAN GENOME SCIENCES, INC.  
9410 KEY WEST AVENUE  
ROCKVILLE, MD 20850  
UNITED STATES OF AMERICA  
  
THE REGENTS OF THE UNIVERSITY OF MICHIGAN  
WOLVERINE TOWER, ROOM 2071  
3003 SOUTH STATE STREET  
ANN ARBOR, MI 48109  
UNITED STATES OF AMERICA
- INVENTORS: NI, JIAN  
ROSEN, CRAIG A.  
DIXIT, VISHVA M.  
GENTZ, REINER L.  
KENNY, JOSEPH J.
- (ii) TITLE OF INVENTION: I-FLICE, A NOVEL INHIBITOR OF TUMOR  
NECROSIS FACTOR RECEPTOR-1 AND CD-95 INDUCED APOPTOSIS
- (iii) NUMBER OF SEQUENCES: 32
- (iv) CORRESPONDENCE ADDRESS:
  - (A) ADDRESSEE: STERNE, KESSLER, GOLDSTEIN & FOX P.L.L.C.
  - (B) STREET: 1100 NEW YORK AVENUE, SUITE 600
  - (C) CITY: WASHINGTON
  - (D) STATE: DC
  - (E) COUNTRY: US
  - (F) ZIP: 20005
- (v) COMPUTER READABLE FORM:
  - (A) MEDIUM TYPE: Floppy disk
  - (B) COMPUTER: IBM PC compatible
  - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
  - (D) SOFTWARE: PatentIn Release #1.0, Version #1.30
- (vi) CURRENT APPLICATION DATA:
  - (A) APPLICATION NUMBER: TO BE ASSIGNED
  - (B) FILING DATE: HERewith
  - (C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:
  - (A) APPLICATION NUMBER: US 60/034,205
  - (B) FILING DATE: 21-JAN-1997
- (vii) PRIOR APPLICATION DATA:
  - (A) APPLICATION NUMBER: US 60/054,800
  - (B) FILING DATE: 05-AUG-1997
- (viii) ATTORNEY/AGENT INFORMATION:
  - (A) NAME: STEFFE, ERIC K.
  - (B) REGISTRATION NUMBER: 36,688
  - (C) REFERENCE/DOCKET NUMBER: 1488.097PC02
- (ix) TELECOMMUNICATION INFORMATION:
  - (A) TELEPHONE: (202) 371-2600
  - (B) TELEFAX: (202) 371-2540

- 68 -

## (2) INFORMATION FOR SEQ ID NO:1:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2034 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: cDNA

## (ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 268..1707

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

CGATCGCCCA GCACCAAGTC CGCTTCAGG CTTTCGGTTT CTTTGCCTCC ATCTTGGGTG	60
CGCCTTCCCG GCGTCTAGGG GAGCGAAGGC TGAGGTGGCA GCGGCAGGAG AGTCCGGCCG	120
CGACAGGACG AACTCCCCCA CTGGAAAGGA TTCTGAAAGA AATGAAGTCA GCCCTCAGAA	180
ATGAAGTTGA CTGCCTGCTG GCTTTCTGTT GACTGGCCCG GAGCTGTACT GCAAGACCCT	240
TGTGAGCTTC CCTAGTCTAA GAGTAGG ATG TCT GCT GAA GTC ATC CAT CAG	291
Met Ser Ala Glu Val Ile His Gln	
1 5	
GTT GAA GAA GCA CTT GAT ACA GAT GAG AAG GAG ATG CTG CTC TTT TTG	339
Val Glu Glu Ala Leu Asp Thr Asp Glu Lys Glu Met Leu Leu Phe Leu	
10 15 20	
TGC CGG GAT GTT GCT ATA GAT GTG GTT CCA CCT AAT GTC AGG GAC CTT	387
Cys Arg Asp Val Ala Ile Asp Val Val Pro Pro Asn Val Arg Asp Leu	
25 30 35 40	
CTG GAT ATT TTA CGG GAA AGA GGT AAG CTG TCT GTC GGG GAC TTG GCT	435
Leu Asp Ile Leu Arg Glu Arg Gly Lys Leu Ser Val Gly Asp Leu Ala	
45 50 55	
GAA CTG CTC TAC AGA GTG AGG CGA TTT GAC CTG CTC AAA CGT ATC TTG	483
Glu Leu Leu Tyr Arg Val Arg Arg Phe Asp Leu Leu Lys Arg Ile Leu	
60 65 70	
AAG ATG GAC AGA AAA GCT GTG GAG ACC CAC CTG CTC AGG AAC CCT CAC	531
Lys Met Asp Arg Lys Ala Val Glu Thr His Leu Leu Arg Asn Pro His	
75 80 85	
CTT GTT TCG GAC TAT AGA GTG CTG ATG GCA GAG ATT GGT GAG GAT TTG	579
Leu Val Ser Asp Tyr Arg Val Leu Met Ala Glu Ile Gly Glu Asp Leu	
90 95 100	
GAT AAA TCT GAT GTG TCC TCA TTA ATT TTC CTC ATG AAG GAT TAC ATG	627
Asp Lys Ser Asp Val Ser Ser Leu Ile Phe Leu Met Lys Asp Tyr Met	
105 110 115 120	
GGC CGA GGC AAG ATA AGC AAG GAG AAG AGT TTC TTG GAC CTT GTG GTT	675
Gly Arg Gly Lys Ile Ser Lys Glu Lys Ser Phe Leu Asp Leu Val Val	
125 130 135	

- 69 -

GAG TTG GAG AAA CTA AAT CTG GTT GCC CCA GAT CAA CTG GAT TTA TTA Glu Leu Glu Lys Leu Asn Leu Val Ala Pro Asp Gln Leu Asp Leu Leu 140 145 150	723
GAA AAA TGC CTA AAG AAC ATC CAC AGA ATA GAC CTG AAG ACA AAA ATC Glu Lys Cys Leu Lys Asn Ile His Arg Ile Asp Leu Lys Thr Lys Ile 155 160 165	771
CAG AAG TAC AAG CAG TCT GTT CAA GGA GCA GGG ACA AGT TAC AGG AAT Gln Lys Tyr Lys Gln Ser Val Gln Gly Ala Gly Thr Ser Tyr Arg Asn 170 175 180	819
GTT CTC CAA GCA GCA ATC CAA AAG AGT CTC AAG GAT CCT TCA AAT AAC Val Leu Gln Ala Ala Ile Gln Lys Ser Leu Lys Asp Pro Ser Asn Asn 185 190 195 200	867
TTC AGG CTC CAT AAT GGG AGA AGT AAA GAA CAA AGA CTT AAG GAA CAG Phe Arg Leu His Asn Gly Arg Ser Lys Glu Gln Arg Leu Lys Glu Gln 205 210 215	915
CTT GGC GCT CAA CAA GAA CCA GTG AAG AAA TCC ATT CAG GAA TCA GAA Leu Gly Ala Gln Gln Glu Pro Val Lys Lys Ser Ile Gln Glu Ser Glu 220 225 230	963
GCT TTT TTG CCT CAG AGC ATA CCT GAA GAG AGA TAC AAG ATG AAG AGC Ala Phe Leu Pro Gln Ser Ile Pro Glu Glu Arg Tyr Lys Met Lys Ser 235 240 245	1011
AAG CCC CTA GGA ATC TGC CTG ATA ATC GAT TGC ATT GGC AAT GAG ACA Lys Pro Leu Gly Ile Cys Leu Ile Ile Asp Cys Ile Gly Asn Glu Thr 250 255 260	1059
GAG CTT CTT CGA GAC ACC TTC ACT TCC CTG GGC TAT GAA GTC CAG AAA Glu Leu Leu Arg Asp Thr Phe Thr Ser Leu Gly Tyr Glu Val Gln Lys 265 270 275 280	1107
TTC TTG CAT CTC AGT ATG CAT GGT ATA TCC CAG ATT CTT GGC CAA TTT Phe Leu His Leu Ser Met His Gly Ile Ser Gln Ile Leu Gly Gln Phe 285 290 295	1155
GCC TGT ATG CCC GAG CAC CGA GAC TAC GAC AGC TTT GTG TGT GTC CTG Ala Cys Met Pro Glu His Arg Asp Tyr Asp Ser Phe Val Cys Val Leu 300 305 310	1203
GTG AGC CGA GGA GGC TCC CAG AGT GTG TAT GGT GTG GAT CAG ACT CAC Val Ser Arg Gly Gly Ser Gln Ser Val Tyr Gly Val Asp Gln Thr His 315 320 325	1251
TCA GGG CTC CCC CTG CAT CAC ATC AGG AGG ATG TTC ATG GGA GAT TCA Ser Gly Leu Pro Leu His His Ile Arg Arg Met Phe Met Gly Asp Ser 330 335 340	1299
TGC CCT TAT CTA GCA GGG AAG CCA AAG ATG TTT TTT ATT CAG AAC TAT Cys Pro Tyr Leu Ala Gly Lys Pro Lys Met Phe Phe Ile Gln Asn Tyr 345 350 355 360	1347
GTG GTG TCA GAG GGC CAG CTG GAG GAC AGC AGC CTC TTG GAG GTG GAT Val Val Ser Glu Gly Gln Leu Glu Asp Ser Ser Leu Leu Glu Val Asp 365 370 375	1395
GGG CCA GCG ATG AAG AAT GTG GAA TTC AAG GCT CAG AAG CGA GGG CTG Gly Pro Ala Met Lys Asn Val Glu Phe Lys Ala Gln Lys Arg Gly Leu 380 385 390	1443

- 70 -

TGC ACA GTT CAC CGA GAA GCT GAC TTC TTC TGG AGC CTG TGT ACT GCG	1491
Cys Thr Val His Arg Glu Ala Asp Phe Phe Trp Ser Leu Cys Thr Ala	
395 400 405	
GAC ATG TCC CTG CTG GAG CAG TCT CAC AGC TCA CCG TCC CTG TAC CTG	1539
Asp Met Ser Leu Leu Glu Gln Ser His Ser Ser Pro Ser Leu Tyr Leu	
410 415 420	
CAG TGC CTC TCC CAG AAA CTG AGA CAA GAA AGA AAA CGC CCA CTC CTG	1587
Gln Cys Leu Ser Gln Lys Leu Arg Gln Glu Arg Lys Arg Pro Leu Leu	
425 430 435 440	
GAT CTT CAC ATT GAA CTC AAT GGC TAC ATG TAT GAT TGG AAC AGC AGA	1635
Asp Leu His Ile Glu Leu Asn Gly Tyr Met Tyr Asp Trp Asn Ser Arg	
445 450 455	
GTT TCT GCC AAG GAG AAA TAT TAT GTC TGG CTG CAG CAC ACT CTG AGA	1683
Val Ser Ala Lys Glu Lys Tyr Tyr Val Trp Leu Gln His Thr Leu Arg	
460 465 470	
AAG AAA CTT ATC CTC TCC TAC ACA TAAGAAACCA AAAGGCTGGG CGTAGTGGCT	1737
Lys Lys Leu Ile Leu Ser Tyr Thr	
475 480	
CGCACCTGTA ATCCCAGCAC TTTGGGAGGC CAAGGAGGGC GGATCACTTC AGGTCAGGAG	1797
TTCGAGACCA GCCTGGCCAA CATGGTAAAC GCTGTCCCTA GTAAGAGTGC AAAAATTAGC	1857
TGGGTGTGGG TGTGGGTACC TGTGTTCCCA GTTACTTGGG AGGCTGAGGT GGGAGGATCT	1917
TTTGAACCCA GGAGTTCAGG GTCATAGCAT GCTGTGATTG TGCCTACGAA TAGCCACTGC	1977
ATACCAACCT GGGCAATATA GCAAGATCCC ATCTTTTAA AAAAAAAAAA AAAAAAA	2034

## (2) INFORMATION FOR SEQ ID NO:2:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 480 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met Ser Ala Glu Val Ile His Gln Val Glu Glu Ala Leu Asp Thr Asp	
1 5 10 15	
Glu Lys Glu Met Leu Leu Phe Leu Cys Arg Asp Val Ala Ile Asp Val	
20 25 30	
Val Pro Pro Asn Val Arg Asp Leu Leu Asp Ile Leu Arg Glu Arg Gly	
35 40 45	
Lys Leu Ser Val Gly Asp Leu Ala Glu Leu Leu Tyr Arg Val Arg Arg	
50 55 60	
Phe Asp Leu Leu Lys Arg Ile Leu Lys Met Asp Arg Lys Ala Val Glu	
65 70 75 80	
Thr His Leu Leu Arg Asn Pro His Leu Val Ser Asp Tyr Arg Val Leu	
85 90 95	

- 71 -

Met Ala Glu Ile Gly Glu Asp Leu Asp Lys Ser Asp Val Ser Ser Leu  
 100 105 110  
 Ile Phe Leu Met Lys Asp Tyr Met Gly Arg Gly Lys Ile Ser Lys Glu  
 115 120 125  
 Lys Ser Phe Leu Asp Leu Val Val Glu Leu Glu Lys Leu Asn Leu Val  
 130 135 140  
 Ala Pro Asp Gln Leu Asp Leu Leu Glu Lys Cys Leu Lys Asn Ile His  
 145 150 155 160  
 Arg Ile Asp Leu Lys Thr Lys Ile Gln Lys Tyr Lys Gln Ser Val Gln  
 165 170 175  
 Gly Ala Gly Thr Ser Tyr Arg Asn Val Leu Gln Ala Ala Ile Gln Lys  
 180 185 190  
 Ser Leu Lys Asp Pro Ser Asn Asn Phe Arg Leu His Asn Gly Arg Ser  
 195 200 205  
 Lys Glu Gln Arg Leu Lys Glu Gln Leu Gly Ala Gln Gln Glu Pro Val  
 210 215 220  
 Lys Lys Ser Ile Gln Glu Ser Glu Ala Phe Leu Pro Gln Ser Ile Pro  
 225 230 235 240  
 Glu Glu Arg Tyr Lys Met Lys Ser Lys Pro Leu Gly Ile Cys Leu Ile  
 245 250 255  
 Ile Asp Cys Ile Gly Asn Glu Thr Glu Leu Leu Arg Asp Thr Phe Thr  
 260 265 270  
 Ser Leu Gly Tyr Glu Val Gln Lys Phe Leu His Leu Ser Met His Gly  
 275 280 285  
 Ile Ser Gln Ile Leu Gly Gln Phe Ala Cys Met Pro Glu His Arg Asp  
 290 295 300  
 Tyr Asp Ser Phe Val Cys Val Leu Val Ser Arg Gly Gly Ser Gln Ser  
 305 310 315 320  
 Val Tyr Gly Val Asp Gln Thr His Ser Gly Leu Pro Leu His His Ile  
 325 330 335  
 Arg Arg Met Phe Met Gly Asp Ser Cys Pro Tyr Leu Ala Gly Lys Pro  
 340 345 350  
 Lys Met Phe Phe Ile Gln Asn Tyr Val Val Ser Glu Gly Gln Leu Glu  
 355 360 365  
 Asp Ser Ser Leu Leu Glu Val Asp Gly Pro Ala Met Lys Asn Val Glu  
 370 375 380  
 Phe Lys Ala Gln Lys Arg Gly Leu Cys Thr Val His Arg Glu Ala Asp  
 385 390 395 400  
 Phe Phe Trp Ser Leu Cys Thr Ala Asp Met Ser Leu Leu Glu Gln Ser  
 405 410 415  
 His Ser Ser Pro Ser Leu Tyr Leu Gln Cys Leu Ser Gln Lys Leu Arg  
 420 425 430

- 72 -

Gln Glu Arg Lys Arg Pro Leu Leu Asp Leu His Ile Glu Leu Asn Gly  
           435                                  440                                  445

Tyr Met Tyr Asp Trp Asn Ser Arg Val Ser Ala Lys Glu Lys Tyr Tyr  
           450                                  455                                  460

Val Trp Leu Gln His Thr Leu Arg Lys Lys Leu Ile Leu Ser Tyr Thr  
           465                                  470                                  475                                  480

## (2) INFORMATION FOR SEQ ID NO:3:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 478 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

Met Asp Phe Ser Arg Asn Leu Tyr Asp Ile Gly Glu Gln Leu Asp Ser  
   1                                  5                                  10                                  15

Glu Asp Leu Ala Ser Leu Lys Phe Leu Ser Leu Asp Tyr Ile Pro Gln  
           20                                  25                                  30

Arg Lys Gln Glu Pro Ile Lys Asp Ala Leu Met Leu Phe Gln Arg Leu  
           35                                  40                                  45

Gln Glu Lys Arg Met Leu Glu Glu Ser Asn Leu Ser Phe Leu Lys Glu  
           50                                  55                                  60

Leu Leu Phe Arg Ile Asn Arg Leu Asp Leu Leu Ile Thr Tyr Leu Asn  
           65                                  70                                  75                                  80

Thr Arg Lys Glu Glu Met Glu Arg Glu Leu Gln Thr Pro Gly Arg Ala  
           85                                  90                                  95

Gln Ile Ser Ala Tyr Arg Val Met Leu Tyr Gln Ile Ser Glu Glu Val  
           100                                  105                                  110

Ser Arg Ser Glu Leu Arg Ser Phe Lys Phe Leu Leu Gln Glu Glu Ile  
           115                                  120                                  125

Ser Lys Cys Lys Leu Asp Asp Asp Met Asn Leu Leu Asp Ile Phe Ile  
           130                                  135                                  140

Glu Met Glu Lys Arg Val Ile Leu Gly Glu Gly Lys Leu Asp Ile Leu  
           145                                  150                                  155                                  160

Lys Arg Val Cys Ala Gln Ile Asn Lys Ser Leu Leu Lys Ile Ile Asn  
           165                                  170                                  175

Asp Tyr Glu Glu Phe Ser Lys Glu Arg Ser Ser Ser Leu Glu Gly Ser  
           180                                  185                                  190

Pro Asp Glu Phe Ser Asn Gly Glu Glu Leu Cys Gly Val Met Thr Ile  
           195                                  200                                  205

- 73 -

```

Ser Asp Ser Pro Arg Glu Gln Asp Ser Glu Ser Gln Thr Leu Asp Lys
210                215                220

Val Tyr Gln Met Lys Ser Lys Pro Arg Gly Tyr Cys Leu Ile Ile Asn
225                230                235                240

Asn His Asn Phe Ala Lys Ala Arg Glu Lys Val Pro Lys Leu His Ser
245                250                255

Ile Arg Asp Arg Asn Gly Thr His Leu Asp Ala Gly Leu Thr Thr Thr
260                265                270

Phe Glu Glu Leu His Phe Glu Ile Lys Pro His Asp Asp Cys Thr Val
275                280                285

Glu Gln Ile Tyr Glu Ile Leu Lys Ile Tyr Gln Leu Met Asp His Ser
290                295                300

Asn Met Asp Cys Phe Ile Cys Cys Ile Leu Ser His Gly Asp Lys Gly
305                310                315                320

Ile Ile Tyr Gly Thr Asp Gly Gln Glu Pro Pro Ile Tyr Glu Leu Thr
325                330                335

Ser Gln Phe Thr Gly Leu Lys Cys Pro Ser Leu Ala Gly Lys Pro Lys
340                345                350

Val Phe Phe Ile Gln Ala Cys Gln Gly Asp Asn Tyr Gln Lys Gly Ile
355                360                365

Pro Val Glu Thr Asp Ser Glu Glu Gln Pro Tyr Leu Glu Met Asp Leu
370                375                380

Ser Ser Pro Gln Thr Arg Tyr Ile Pro Asp Glu Ala Asp Phe Leu Leu
385                390                395                400

Gly Met Ala Thr Val Asn Asn Cys Val Ser Tyr Arg Asn Pro Ala Glu
405                410                415

Gly Thr Trp Tyr Ile Gln Ser Leu Cys Gln Ser Leu Arg Glu Arg Cys
420                425                430

Pro Arg Gly Asp Asp Ile Leu Thr Ile Leu Thr Glu Val Asn Tyr Glu
435                440                445

Val Ser Asn Lys Asp Asp Lys Lys Asn Met Gly Lys Gln Met Pro Gln
450                455                460

Pro Thr Phe Thr Leu Arg Lys Lys Leu Val Phe Pro Ser Asp
465                470                475

```

## (2) INFORMATION FOR SEQ ID NO:4:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 479 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

- 74 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Met	Lys	Ser	Gln	Gly	Gln	His	Trp	Tyr	Ser	Ser	Ser	Asp	Lys	Asn	Cys	1	5	10	15
Lys	Val	Ser	Phe	Arg	Glu	Lys	Leu	Leu	Ile	Ile	Asp	Ser	Asn	Leu	Gly	20	25	30	
Val	Gln	Asp	Val	Glu	Asn	Leu	Lys	Phe	Leu	Cys	Ile	Gly	Leu	Val	Pro	35	40	45	
Asn	Lys	Lys	Leu	Glu	Lys	Ser	Ser	Ser	Ala	Ser	Asp	Val	Phe	Glu	His	50	55	60	
Leu	Leu	Ala	Glu	Asp	Leu	Leu	Ser	Glu	Glu	Asp	Pro	Phe	Phe	Leu	Ala	65	70	75	80
Glu	Leu	Leu	Tyr	Ile	Ile	Arg	Gln	Lys	Lys	Leu	Leu	Gln	His	Leu	Asn	85	90	95	
Cys	Thr	Lys	Glu	Glu	Val	Glu	Arg	Leu	Leu	Pro	Thr	Arg	Gln	Arg	Val	100	105	110	
Ser	Leu	Phe	Arg	Asn	Leu	Leu	Tyr	Glu	Leu	Ser	Glu	Gly	Ile	Asp	Ser	115	120	125	
Glu	Asn	Leu	Lys	Asp	Met	Ile	Phe	Leu	Leu	Lys	Asp	Ser	Leu	Pro	Lys	130	135	140	
Thr	Glu	Met	Thr	Ser	Leu	Ser	Phe	Leu	Ala	Phe	Leu	Glu	Lys	Gln	Gly	145	150	155	160
Lys	Ile	Asp	Glu	Asp	Asn	Leu	Thr	Cys	Leu	Glu	Asp	Leu	Cys	Lys	Thr	165	170	175	
Val	Val	Pro	Lys	Leu	Leu	Arg	Asn	Ile	Glu	Lys	Tyr	Lys	Arg	Glu	Lys	180	185	190	
Ala	Ile	Gln	Ile	Val	Thr	Pro	Pro	Val	Asp	Lys	Glu	Ala	Glu	Ser	Tyr	195	200	205	
Gln	Gly	Glu	Glu	Glu	Leu	Val	Ser	Gln	Thr	Asp	Val	Lys	Thr	Phe	Leu	210	215	220	
Glu	Ala	Leu	Pro	Arg	Ala	Ala	Val	Tyr	Arg	Met	Asn	Arg	Asn	His	Arg	225	230	235	240
Gly	Leu	Cys	Val	Ile	Val	Asn	Asn	His	Ser	Phe	Thr	Ser	Leu	Lys	Asp	245	250	255	
Arg	Gln	Gly	Thr	His	Lys	Asp	Ala	Glu	Ile	Leu	Ser	His	Val	Phe	Gln	260	265	270	
Trp	Leu	Gly	Phe	Thr	Val	His	Ile	His	Asn	Asn	Val	Thr	Lys	Val	Glu	275	280	285	
Met	Glu	Met	Val	Leu	Gln	Lys	Gln	Lys	Cys	Asn	Pro	Ala	His	Ala	Asp	290	295	300	
Gly	Asp	Cys	Phe	Val	Phe	Cys	Ile	Leu	Thr	His	Gly	Arg	Phe	Gly	Ala	305	310	315	320
Val	Tyr	Ser	Ser	Asp	Glu	Ala	Leu	Ile	Pro	Ile	Arg	Glu	Ile	Met	Ser				



- 75 -

	325		330		335
His Phe Thr	Ala Leu Gln Cys Pro	Arg Leu Ala Glu Lys	Pro Lys Leu		
	340	345	350		
Phe Phe Ile	Gln Ala Cys Gln Gly	Glu Glu Ile Gln	Pro Ser Val Ser		
	355	360	365		
Ile Glu Ala	Asp Ala Leu Asn Pro	Glu Gln Ala Pro	Thr Ser Leu Gln		
	370	375	380		
Asp Ser Ile	Pro Ala Glu Ala Asp	Phe Leu Leu Gly	Leu Ala Thr Val		
	385	390	395	400	
Pro Gly Tyr	Val Ser Phe Arg His	Val Glu Glu Gly	Ser Trp Tyr Ile		
	405	410	415		
Gln Ser Leu	Cys Asn His Leu Lys	Lys Leu Val Pro	Arg His Glu Asp		
	420	425	430		
Ile Leu Ser	Ile Leu Thr Ala Val	Asn Asp Asp Val	Ser Arg Arg Val		
	435	440	445		
Asp Lys Gln	Gly Thr Lys Lys Gln	Met Pro Gln Pro	Ala Phe Thr Leu		
	450	455	460		
Arg Lys Lys	Leu Val Phe Pro	Val Pro Leu Asp	Ala Leu Ser Ile		
	465	470	475		

## (2) INFORMATION FOR SEQ ID NO:5:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2597 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: cDNA

## (ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 304..1347

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

GCGAGCTTGC AGCCTCACCG ACGAGTCTCA ACTAAAAGGG ACTCCCGGAG CTAGGGGTGG	60
GGACTCGGCC TCACACAGTG ATTGCCGGCT ATTGGACTTT TGTCCAGTGA CAGCTGAGAC	120
AACAAGGACC ACGGGAGGAG GTGTAGGAGA GAAGCGCCGC GAACAGGCAT CGCCCAGCAC	180
CAAGTCCGCT TCCAGGCTTT CGGTTTCTTT GCCTCCATCT TGGGTGCGCC TTCCCGGCGT	240
CTAGGGGAGC GAAGGCTGAG GTGGCAGCGG CAGGAGAGTC CGGCCGCGAC AGGACGAGTG	300
CTG ATG GCA GAG ATT GGT GAG GAT TTG GAT AAA TCT GAT GTG TCC TCA	348
Met Ala Glu Ile Gly Glu Asp Leu Asp Lys Ser Asp Val Ser Ser	
5 10 15	
TTA ATT TTC CTC ATG AAG GAT TAC ATG GGC CGA GGC AAG ATA AGC AAG	396
Leu Ile Phe Leu Met Lys Asp Tyr Met Gly Arg Gly Lys Ile Ser Lys	

- 76 -

20					25					30							
GAG	AAG	AGT	TTC	TTG	GAC	CTT	GTG	GTT	GAG	TTG	GAG	AAA	CTA	AAT	CTG	444	
Glu	Lys	Ser	Phe	Leu	Asp	Leu	Val	Val	Glu	Leu	Glu	Lys	Leu	Asn	Leu		
35					40					45							
GTT	GCC	CCA	GAT	CAA	CTG	GAT	TTA	TTA	GAA	AAA	TGC	CTA	AAG	AAC	ATC	492	
Val	Ala	Pro	Asp	Gln	Leu	Asp	Leu	Leu	Glu	Lys	Cys	Leu	Lys	Asn	Ile		
50					55					60							
CAC	AGA	ATA	GAC	CTG	AAG	ACA	AAA	ATC	CAG	AAG	TAC	AAG	CAG	TCT	GTT	540	
His	Arg	Ile	Asp	Leu	Lys	Thr	Lys	Ile	Gln	Lys	Tyr	Lys	Gln	Ser	Val		
65					70					75							
CAA	GGA	GCA	GGG	ACA	AGT	TAC	AGG	AAT	GTT	CTC	CAA	GCA	GCA	ATC	CAA	588	
Gln	Gly	Ala	Gly	Thr	Ser	Tyr	Arg	Asn	Val	Leu	Gln	Ala	Ala	Ile	Gln		
80					85					90					95		
AAG	AGT	CTC	AAG	GAT	CCT	TCA	AAT	AAC	TTC	AGG	GAA	GAA	CCA	GTG	AAG	636	
Lys	Ser	Leu	Lys	Asp	Pro	Ser	Asn	Asn	Phe	Arg	Glu	Glu	Pro	Val	Lys		
100					105					110							
AAA	TCC	ATT	CAG	GAA	TCA	GAA	GCT	TTT	TTG	CCT	CAG	AGC	ATA	CCT	GAA	684	
Lys	Ser	Ile	Gln	Glu	Ser	Glu	Ala	Phe	Leu	Pro	Gln	Ser	Ile	Pro	Glu		
115					120					125							
GAG	AGA	TAC	AAG	ATG	AAG	AGC	AAG	CCC	CTA	GGA	ATC	TGC	CTG	ATA	ATC	732	
Glu	Arg	Tyr	Lys	Met	Lys	Ser	Lys	Pro	Leu	Gly	Ile	Cys	Leu	Ile	Ile		
130					135					140							
GAT	TGC	ATT	GGC	AAT	GAG	ACA	GAG	CTT	CTT	CGA	GAC	ACC	TTC	ACT	TCC	780	
Asp	Cys	Ile	Gly	Asn	Glu	Thr	Glu	Leu	Leu	Arg	Asp	Thr	Phe	Thr	Ser		
145					150					155							
CTG	GGC	TAT	GAA	GTC	CAG	AAA	TTC	TTG	CAT	CTC	AGT	ATG	CAT	GGT	ATA	828	
Leu	Gly	Tyr	Glu	Val	Gln	Lys	Phe	Leu	His	Leu	Ser	Met	His	Gly	Ile		
160					165					170					175		
TCC	CAG	ATT	CTT	GGC	CAA	TTT	GCC	TGT	ATG	CCC	GAG	CAC	CGA	GAC	TAC	876	
Ser	Gln	Ile	Leu	Gly	Gln	Phe	Ala	Cys	Met	Pro	Glu	His	Arg	Asp	Tyr		
180					185					190							
GAC	AGC	TTT	GTG	TGT	GTC	CTG	GTG	AGC	CGA	GGA	GGC	TCC	CAG	AGT	GTG	924	
Asp	Ser	Phe	Val	Cys	Val	Leu	Val	Ser	Arg	Gly	Gly	Ser	Gln	Ser	Val		
195					200					205							
TAT	GGT	GTG	GAT	CAG	ACT	CAC	TCA	GGG	CTC	CCC	CTG	CAT	CAC	ATC	AGG	972	
Tyr	Gly	Val	Asp	Gln	Thr	His	Ser	Gly	Leu	Pro	Leu	His	His	Ile	Arg		
210					215					220							
AGG	ATG	TTC	ATG	GGA	GAT	TCA	TGC	CCT	TAT	CTA	GCA	GGG	AAG	CCA	AAG	1020	
Arg	Met	Phe	Met	Gly	Asp	Ser	Cys	Pro	Tyr	Leu	Ala	Gly	Lys	Pro	Lys		
225					230					235							
ATG	TTT	TTT	ATT	CAG	AAC	TAT	GTG	GTG	TCA	GAC	GGC	CAG	CTG	GAG	GAC	1068	
Met	Phe	Phe	Ile	Gln	Asn	Tyr	Val	Val	Ser	Asp	Gly	Gln	Leu	Glu	Asp		
240					245					250					255		
AGC	AGC	CTC	TTG	GAG	GTG	GAT	GGG	CCA	GCG	ATG	AAG	AAT	GTG	GAA	TTC	1116	
Ser	Ser	Leu	Leu	Glu	Val	Asp	Gly	Pro	Ala	Met	Lys	Asn	Val	Glu	Phe		
260					265					270							
AAG	GCT	CAG	AAG	CGA	GGG	CTG	TGC	ACA	GTT	CAC	CGA	GAA	GCT	GAC	TTC	1164	

- 77 -

Lys	Ala	Gln	Lys	Arg	Gly	Leu	Cys	Thr	Val	His	Arg	Glu	Ala	Asp	Phe	
			275					280					285			
TTC	TGG	AGC	CTG	TGT	ACT	GCG	GAC	ATG	TCC	CTG	CTG	GAG	CAG	TCT	CAC	1212
Phe	Trp	Ser	Leu	Cys	Thr	Ala	Asp	Met	Ser	Leu	Leu	Glu	Gln	Ser	His	
		290					295					300				
AGC	TCA	CCG	TCC	CTG	TAC	CTG	CAG	TGC	CTC	TCC	CAG	AAA	CTG	AGA	CAA	1260
Ser	Ser	Pro	Ser	Leu	Tyr	Leu	Gln	Cys	Leu	Ser	Gln	Lys	Leu	Arg	Gln	
	305					310					315					
GAA	AGG	GGG	ACA	ATT	CCC	GGA	AGT	GGA	ATT	ACA	GAG	TCA	AAG	GAC	ATG	1308
Glu	Arg	Gly	Thr	Ile	Pro	Gly	Ser	Gly	Ile	Thr	Glu	Ser	Lys	Asp	Met	
320				325					330						335	
CAT	TTT	TCA	AGC	CTC	GGA	TGC	ATC	TTA	CTA	GAT	GTC	CTA	TAGGATGGTC			1357
His	Phe	Ser	Ser	Leu	Gly	Cys	Ile	Leu	Leu	Asp	Val	Leu				
			340					345								
ATATCAGCTT	TATAGGAGAG	TAGCTGTGTC	CCTGAATTCT	CCCTGACACT	GCATGCTCTT											1417
ATATTTCCCTC	AAGTTTTGAC	AATTTGATAG	GTGAAAAGTG	GSTATCTGACT	GTTTCAGATCT											1477
GGAAGGCTTT	GTTATATAAA	CATTTTTTTA	ATGTTTATTG	GCAAGAATAC	TTTTCTAAGA											1537
GAAACATCAG	TGAGCTGGTT	TCCATTTAAG	CTGAATGAAG	CCACAATGTA	CCTCAAGTAT											1597
AAGATTAAC	GGCCTTTTTT	AGTTGCACTC	TAATTACAAT	TTAGAATGAT	GTTTCTGAGC											1657
CACCTGTCAA	ATGCATTCTG	GGCTGTACCT	CTGCGTACCC	CAGGAATAAA	TCTCATGGCC											1717
TTCTTTACCT	GGCCTCCTTA	GTGGTGGCCC	AGCAGGAAGC	GGGGGTTAGA	GCAGGAGCCA											1777
CTCAGCCTTC	CAAGATAGAT	ACTCCATGGG	CCGGTGGTAT	TACTGGCCTT	TTGAGCCCAT											1837
CCCCATTTGC	ATAGATGATC	CACGTGGGTT	ATCATCTGGC	TGGTATGTTT	CCAGAGTGAA											1897
ACTCAGCAGC	CCCTTGAGGG	AGGGGATGGT	GGCCATCAGG	CCAGAGTATT	GCAAGTTAGT											1957
TTGGATCATT	TGCTAAGCAG	CTTGTGGTGC	CTTCAGAAAG	GAACAGTTTC	AAAGAACTTT											2017
CACATCTGTT	GGCTCATTTT	GCCCTAATGA	CAGTCTTCTC	TTTGATATTT	GCATGGCATT											2077
AAATTTTGCC	TTTCTTGTTT	TCTCCAGAAA	ACGCCCACTC	CTGGATCTTC	ACATTGAACT											2137
CAATGGCTAC	ATGTATGATT	GGAACAGCAG	AGTTTCTGCC	AAGGAGAAAT	ATTATGTCTG											2197
GCTGCAGCAC	ACTCTGAGAA	AGAACTTAT	CTCTCCTACA	CATAAGAAAC	CAAAAGGCTG											2257
GGCGTAGTGG	CTCGCACCTG	TGATCCCAGC	ACTTTGGGAG	GCCGAGGAGG	GCGGATCACT											2317
TCAGGTCGGG	AGTTCGAGAC	CAGCCTGGCC	AGCATGTGAA	CGCTGTCCCT	AGTAGAAGTG											2377
CAAAAATTGG	CTGGTGTGGG	TGTGGGTACC	CTGTATTCCC	AGTTGCTTGG	GGGGCTGAGG											2437
TGGGAGGATC	TTTTGACCCC	AGGAGTTCAG	GGTCATAGCA	TGCTGTGATT	GTGCCTACGA											2497
ATAGCCACTG	CATACCAACC	TGGGCAATAT	AGCAAGATCC	CATCTCTTTA	AAAAAAAAAA											2557
AAAAAGGACA	GGAACATCT	TAAAAAAAAA	AAAAAAAAAA													2597

(2) INFORMATION FOR SEQ ID NO:6:

- 78 -

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 348 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

```

Met Ala Glu Ile Gly Glu Asp Leu Asp Lys Ser Asp Val Ser Ser Leu
 1           5           10           15
Ile Phe Leu Met Lys Asp Tyr Met Gly Arg Gly Lys Ile Ser Lys Glu
 20           25           30
Lys Ser Phe Leu Asp Leu Val Val Glu Leu Glu Lys Leu Asn Leu Val
 35           40           45
Ala Pro Asp Gln Leu Asp Leu Leu Glu Lys Cys Leu Lys Asn Ile His
 50           55           60
Arg Ile Asp Leu Lys Thr Lys Ile Gln Lys Tyr Lys Gln Ser Val Gln
 65           70           75           80
Gly Ala Gly Thr Ser Tyr Arg Asn Val Leu Gln Ala Ala Ile Gln Lys
 85           90           95
Ser Leu Lys Asp Pro Ser Asn Asn Phe Arg Glu Glu Pro Val Lys Lys
 100          105          110
Ser Ile Gln Glu Ser Glu Ala Phe Leu Pro Gln Ser Ile Pro Glu Glu
 115          120          125
Arg Tyr Lys Met Lys Ser Lys Pro Leu Gly Ile Cys Leu Ile Ile Asp
 130          135          140
Cys Ile Gly Asn Glu Thr Glu Leu Leu Arg Asp Thr Phe Thr Ser Leu
 145          150          155          160
Gly Tyr Glu Val Gln Lys Phe Leu His Leu Ser Met His Gly Ile Ser
 165          170          175
Gln Ile Leu Gly Gln Phe Ala Cys Met Pro Glu His Arg Asp Tyr Asp
 180          185          190
Ser Phe Val Cys Val Leu Val Ser Arg Gly Gly Ser Gln Ser Val Tyr
 195          200          205
Gly Val Asp Gln Thr His Ser Gly Leu Pro Leu His His Ile Arg Arg
 210          215          220
Met Phe Met Gly Asp Ser Cys Pro Tyr Leu Ala Gly Lys Pro Lys Met
 225          230          235          240
Phe Phe Ile Gln Asn Tyr Val Val Ser Asp Gly Gln Leu Glu Asp Ser
 245          250          255
Ser Leu Leu Glu Val Asp Gly Pro Ala Met Lys Asn Val Glu Phe Lys
 260          265          270
Ala Gln Lys Arg Gly Leu Cys Thr Val His Arg Glu Ala Asp Phe Phe
 275          280          285

```

- 79 -

Trp Ser Leu Cys Thr Ala Asp Met Ser Leu Leu Glu Gln Ser His Ser  
 290 295 300

Ser Pro Ser Leu Tyr Leu Gln Cys Leu Ser Gln Lys Leu Arg Gln Glu  
 305 310 315 320

Arg Gly Thr Ile Pro Gly Ser Gly Ile Thr Glu Ser Lys Asp Met His  
 325 330 335

Phe Ser Ser Leu Gly Cys Ile Leu Leu Asp Val Leu  
 340 345

## (2) INFORMATION FOR SEQ ID NO:7:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 26 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

CGCCCATGGC TGAAGTCATC CATCAG

26

## (2) INFORMATION FOR SEQ ID NO:8:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 27 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

CGCAAGCTTG TGCTGGGATT ACAGGTG

27

## (2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 26 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

CGCCCATGGA GATTGGTGAG GATTTG

26

- 80 -

## (2) INFORMATION FOR SEQ ID NO:10:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 26 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

CGCAAGCTTA GAGCATGCAG TGTCAG

26

## (2) INFORMATION FOR SEQ ID NO:11:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 33 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

CGCGGATCCG CCATCATGTC TGCTGAAGTC ATC

33

## (2) INFORMATION FOR SEQ ID NO:12:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 27 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

CGCGGTACCG TGCTGGGATT ACAGGTG

27

## (2) INFORMATION FOR SEQ ID NO:13:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 33 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- 81 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

CGCGGATCCG CCATCATGGC AGAGATTGGT GAG

33

(2) INFORMATION FOR SEQ ID NO:14:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 26 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

CGCGGTACCA GAGCATGCAG TGTCAG

26

(2) INFORMATION FOR SEQ ID NO:15:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 33 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

CGCCCCGGGG CCATCATGTC TGCTGAAGTC ATC

33

(2) INFORMATION FOR SEQ ID NO:16:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 57 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

CGCTCTAGAT CAAGCGTAGT CTGGGACGTC GTATGGGTAG TGCTGGGATT ACAGGTG

57

(2) INFORMATION FOR SEQ ID NO:17:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 33 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

- 82 -

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

CGCGGATCCG CCATCATGTC TGCTGAAGTC ATC

33

(2) INFORMATION FOR SEQ ID NO:18:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 27 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

CGCGGTACCG TGCTGGGATT ACAGGTG

27

(2) INFORMATION FOR SEQ ID NO:19:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 33 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

CGCGGATCCG CCATCATGGC AGAGATTGGT GAG

33

(2) INFORMATION FOR SEQ ID NO:20:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 56 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

CGCTCTAGAT CAAGCGTAGT CTGGGACGTC GTATGGGTAA GAGCATGCAG TGTCAG

56

(2) INFORMATION FOR SEQ ID NO:21:

- (i) SEQUENCE CHARACTERISTICS:



- 83 -

- (A) LENGTH: 33 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

CGCGGATCCG CCATCATGGC AGAGATTGGT GAG

33

(2) INFORMATION FOR SEQ ID NO:22:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 26 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

CGCGGTACCA GAGCATGCAG TGTCAG

26

(2) INFORMATION FOR SEQ ID NO:23:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 414 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

AATTCGGCAC GAGGGNGGAC TTGGCTGAAC TGCTCTACAG AGTGAGGCGA TTTGACCTGC	60
TCAAACGTAT CTTGAAGATG GACAGAAAAG CTGTGGAGAC CCACCTGCTC AGGAACCCTC	120
ACCTTGTTTC GGACTATAGA GTGCTGATGG CAGAGATTGG TGAGGATTTG GATAAATCTG	180
ATGTGTCCTC ATTAATTTTC CTCATGAAGG ATTACATGGG CCGAGGCAAG ATAAGCAAGG	240
AGAAGAGTTT CTTGGGACCT TGGTGGTTGA GTTGGGAGAA ACTAAATCTG GTTTGCCCCA	300
GATCAACTNG GGATTNTTA GGAAAAATGC CTAAAGAACA TNCACAGGAT AGACCTGNAG	360
ACAAAANTCC AGNAGTACAN GCAGTNTGTT CAGGGAGCAG GGACAATTNC AGGA	414

(2) INFORMATION FOR SEQ ID NO:24:

- (i) SEQUENCE CHARACTERISTICS:

- 84 -

- (A) LENGTH: 393 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

TGCCAAGGAG AAATATTATG TCTGGCTGCA GCACACTCTG NGAAAGAAAC TTATCCTCTC	60
CTACACATAA GAAACCNAAG GGCTGGGCGT AGTGGCTCAC GCCTGTNAAT CCCAGCACTT	120
TGGGAGGCCA AGGAGGGCAG ATCACTTCAG GTCAGGAGTT CGAGACCAGC CTGGCCAACA	180
TGGTAAACGC TGTCCCTAGT AAAANTACAA AANTTAGCTG GGTGTGGGTG TGGGTACCTG	240
TGTTCCCAGT TACTTGGGAG GCTGAGGTGG GAGGATCTTT TGGAAACCCAG GAGTTTCAGG	300
GTCATAGCAT GCTGTGNTTG TGCCCTNACG AATTAGCCAC TGCATTACCA ACCTGGGGCA	360
ATNTTAGGCA AGATCCCATN TTTTTTAAAA AAA	393

(2) INFORMATION FOR SEQ ID NO:25:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 309 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

TGGATCTTCA CATTGAACTC AATGGCTACA TGTATGATTG GAACAGCAGA GTTCTGCCA	60
AGGAGAAATA TTATGTCTGG CTGCAGCACA CTCTGAGAAA GAAACTTATC CTCTCCTACA	120
CATAAGAAAC CAAAAGGCTG GCGTAGTGG CTCACGCCTG TGATCCCAGC ACTTTGGGAG	180
GCCGGGGAGG GCAGATCACT TCAGGTCAGG AGTTCGAGAC CGGCCTGGNC AACATGGTAG	240
ACGCTGTCCC TAGTAAAAAT GCAAAAGTTG GCTGGGTGTG GGTGTNGGTA CCTGTGTTCC	300
CAGTTGCTT	309

(2) INFORMATION FOR SEQ ID NO:26:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 500 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- 85 -

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

AATTCGGCAG AGCTCACTCA GGGCTCCCC TGCATCACAT CAGGAGGATG TTCATGGGAG	60
ATTCATGCCC TTATCTAGCA GGAAGCCAA AGATGTTTTT TATTCAGAAC TATGTGNTGT	120
CAGAGGGCCA GCTGGAGGAC AGCAGCCTCT TGGAGGTGGA TGGGCCAGCN ATGAAGAATG	180
TGGAATTCAA GGCTCAGAAG CGAGGGCTGT GCACAGTTCA CCGAGNAAGC TGACTTCTTC	240
TGGAGCCTGT GTAATGCGGA CATGTCCCTG CTTGGAGCAA TCTTCANAGG TTCANCGTCC	300
CTGTNACCTG CATGCCTTTT CCCAGAACT GNGACAAGNA AGAAAACGNC CANTNCTGGG	360
GNTNTTTCAC ATTGGAATC AATGGTTACA ANTTATGNTT GGGGNCAACA ANTTTTTTGC	420
CAAGGGGGAA TTTTTGTTT TGGGNTGNAG AAAAATTNG GGAAAGAANT TTTCCCTTN	480
CNNNAAATTA GGNACCCAAA	500

## (2) INFORMATION FOR SEQ ID NO:27:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 324 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: cDNA

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

ATTCTGAAAA AGAATGTGGG GTTTCCTTGC AGATGAGTTC ATCTGTTGTT TCATTTTCCTT	60
TACAATAACT CCCCCACTGG AAAGGATTCT GAAAGAAATG AAGTCAGCCC TCAGAAATGA	120
AGTTGACTGC CTGCTGGCTT TCTGTTGACT GGCCTGGAGC TGTACTGCAA GACCCTTGTC	180
AGCTTCCCTA GTCTAAGAGT AGGATGTCTG CTGAAGTCAT CCATCAGGTT GAAGAAGCAC	240
TTGATACAGA TGAGAAGGAG ATGCTGCTCT TTTTGTGCCG GGATGTTGCT ATAGATGTGG	300
TTCCACCTAA TGTCAGGGAC CTTC	324

## (2) INFORMATION FOR SEQ ID NO:28:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 389 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: cDNA

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

- 86 -

NAATTCGGCA GAGANAAGAG TCTCAAGGAT CCTTCAAATA ACTTCAGGCT CCATAATGGG	60
AGAAGTAAAG AACAAAGACT TAAGGAACAG CTTGGCGCTC AACAGAACC AGTGNAAGAA	120
ATCCATTGAG GAATCAGAAG CTTTTTTGCC TCAGAGCATA CCTGAAGAGA GATACAAGAT	180
GAAGAGCAAG CCCCTAGGGA ATCTGCCTGA TAAATCGATT GCATTGGCAA TGAGGACAGA	240
GCTTCTTCGG GGACACCTTC ACTTCCCTGG GCTTATGAAG TNCCAGGAAA TTCTTGCATC	300
TCAGTATGCA TGGTATTNTC CCAGATTTTT TGGGNCCAAT TTGCCCGTTA TGNCCNGGGC	360
ANCNGGGGAT TTANGACAAT TTTGTGGTG	389

## (2) INFORMATION FOR SEQ ID NO:29:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 308 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

ATTCTGAAAA AGANTGNNGG GTTTCCTTGC AGATGAGTTC ATCTNTTGTT TCATTTTCCTT	60
TACAATAACT CCCCCACTGG AAAGGATTCT GAAAGNAATG AAGTCAGCCC TCAGAAATGA	120
AGTTGNCTGC CTGCTGGCTT TCTGTTGACT GGCCTGGAGC TGTAAGTCAA GACCCTTGTTG	180
AGCTTCCCTA GTCTAAGAGT AGGATGTCTG CTGAAGTCAT CCATCAGGTT GAAGAAGCAC	240
TTGATACAGA TGAGAAGGAG ATGCTGCTCT TTTTGTGCCG GGATGTTTGC TATAGATGTG	300
GTTCCACC	308

## (2) INFORMATION FOR SEQ ID NO:30:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 297 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

ATTCTGAAAA AGAATGTGGG GTTTCCTNGC AGATGAGTTC ATCTGTTGTT TCATTTTCCTT	60
TACAATAACT CCCCCACTGG AAAGGATTCT GAAAGAAATG AAGTCAGCCC TCAGAAATGA	120
AGTTGACTGC CTGCTGGCTT TCTGTTGACT GGCCTGGAGC TGTAAGTCAA GACCCTTGTTG	180
AGCTTCCCTA GTCTAAGAGT AGGATGTCTG CTGAAGTCAT CCATCAGGTT GAAGAAGCAC	240

- 87 -

TTGATACAGA TGAGAAGGAG ATGCTGCTCT TTTTGTGCC GGGATGTTGC TATAGAT 297

(2) INFORMATION FOR SEQ ID NO:31:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 348 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

TATAGGATGG TCATATCAGC TTTATAGGAG AGTAGCTGTG TCCCTGAATT CTCCCTGACA	60
CTGCATGCTC TTATATTTCC TCAAGTTTTG ACAATTTGAT AGGTGAAAAG TGGTATCTGA	120
CTGTTTCAGAT CTGGAAGGCT TTGTTATATA AACATTTTTT TAATGTTTAT TGGCAAGAAT	180
ACTTTTCTAA GAGAAACATC AGTGAGCTGG TTTCCATTTA AGCTGAATGA AGCCACAATG	240
TACCTCAAGT ATAAGGTTAA CTGGCCTTTT TTCAGTTGCA CTCTAATTAC AATTTAGAAT	300
GATGTTTCTG AGCCACCTGT CAAATGCATT CTGGGGCTGT ACCTCTTG	348

(2) INFORMATION FOR SEQ ID NO:32:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 333 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

TATAGGATGG TCATATCAGC TTTATAGGAG AGTAGCTGTG TCCCTGANTT CTCCCTGACA	60
CTGCATGCTC TTATATTTCC TCAAGTTTTG ACAATTTGAT AGGTGAAAAG TGGTATCTGA	120
CTGTNCAGAT CTGGAAGGCT TTGTTATATA AACATTTTTT TAATGTTTAT TGGCAAGAAT	180
ACTTTTCTAA GAGAAACATC AGTGAGCTGG TTTCCATTTA AGCTGAATGA AGCCACAATG	240
TACCTCANGT ATAAGGATTA ACTGGCCTTT TTCCAGTTGC ACTCTAATTA CAATTTTAGA	300
ATGATGTTCN GAGGCCACCT GTCAAATGCA TTC	333

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM  
(PCT Rule 13bis)


A. The indications made below relate to the microorganism referred to in the description on page <u>3</u> , line <u>28</u> .	
<b>B. IDENTIFICATION OF DEPOSIT</b> <span style="float: right;">Further deposits are identified on an additional sheet <input type="checkbox"/></span>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country)  12301 Parklawn Drive Rockville, Maryland 20852 United States of America	
Date of deposit May 15, 1997	Accession Number ATCC 209038
<b>C. ADDITIONAL INDICATIONS</b> (leave blank if not applicable) <span style="float: right;">This information is continued on an additional sheet <input type="checkbox"/></span>	
DNA Plasmid 1631454	
<b>D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE</b> (if the indications are not for all designated States)	
<b>E. SEPARATE FURNISHING OF INDICATIONS</b> (leave blank if not applicable)	
The indications listed below will be submitted to the international Bureau later (specify the general nature of the indications. e.g., "Accession Number of Deposit")	

For receiving Office use only
<input checked="" type="checkbox"/> This sheet was received with the international application
Authorized officer <b>Sonya Barnes</b> <b>PCT International Division</b>

For International Bureau use only
<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM  
(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>4</u> , line <u>5</u> .	
<b>B. IDENTIFICATION OF DEPOSIT</b> <span style="float: right;">Further deposits are identified on an additional sheet <input type="checkbox"/></span>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country)  12301 Parklawn Drive Rockville, Maryland 20852 United States of America	
Date of deposit May 15, 1997	Accession Number ATCC 209041
<b>C. ADDITIONAL INDICATIONS</b> (leave blank if not applicable) <span style="float: right;">This information is continued on an additional sheet <input type="checkbox"/></span>	
DNA Plasmid 1789630	
<b>D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE</b> (if the indications are not for all designated States)	
<b>E. SEPARATE FURNISHING OF INDICATIONS</b> (leave blank if not applicable)	
The indications listed below will be submitted to the international Bureau later (specify the general nature of the indications, e.g., "Accession Number of Deposit")	

For receiving Office use only
<input checked="" type="checkbox"/> This sheet was received with the international application
Authorized officer  PCT International Division

For International Bureau use only
<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer

***What Is Claimed Is:***

1. An isolated nucleic acid molecule comprising a polynucleotide having a nucleotide sequence at least 95% identical to a sequence selected from the group consisting of:

5 (a) a nucleotide sequence encoding a polypeptide comprising amino acids from about 1 to about 480 in SEQ ID NO:2;

(b) a nucleotide sequence encoding a polypeptide comprising amino acids from about 2 to about 480 in SEQ ID NO:2;

10 (c) a nucleotide sequence encoding a polypeptide comprising amino acids from about 1 to about 348 in SEQ ID NO:6;

(d) a nucleotide sequence encoding a polypeptide comprising amino acids from about 2 to about 348 in SEQ ID NO:6;

15 (e) a nucleotide sequence encoding a polypeptide having the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. 209038;

(f) a nucleotide sequence encoding a polypeptide having the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. 209041; and

20 (g) a nucleotide sequence complementary to any of the nucleotide sequences in (a), (b), (c), (d), (e), or (f).

2. An isolated nucleic acid molecule comprising a polynucleotide which hybridizes under stringent hybridization conditions to a polynucleotide having a nucleotide sequence identical to a nucleotide sequence in (a), (b), (c), (d), (e), (f), or (g) of claim 1 wherein said polynucleotide which hybridizes does not hybridize under stringent hybridization conditions to a polynucleotide having a nucleotide sequence consisting of only A residues or of only T residues.

25



3. A method for making a recombinant vector comprising inserting an isolated nucleic acid molecule of claim 1 into a vector.

4. A recombinant vector produced by the method of claim 3.

5. A method of making a recombinant host cell comprising introducing the recombinant vector of claim 4 into a host cell.

6. A recombinant host cell produced by the method of claim 5.

7. A recombinant method for producing an I-FLICE-1 or I-FLICE-2 polypeptide, comprising culturing the recombinant host cell of claim 6 under conditions such that said polypeptide is expressed and recovering said polypeptide.

8. An isolated nucleic acid molecule comprising a polynucleotide having a nucleotide sequence at least 95% identical to a sequence selected from the group consisting of:

(a) a nucleotide sequence encoding a polypeptide comprising amino acids from about 1 to about 75 in SEQ ID NO:2;

(b) a nucleotide sequence encoding a polypeptide comprising amino acids from about 91 to about 171 in SEQ ID NO:2;

(c) a nucleotide sequence encoding a polypeptide comprising amino acids from about 172 to about 375 in SEQ ID NO:2;

(d) a nucleotide sequence encoding a polypeptide comprising amino acids from about 376 to about 480 in SEQ ID NO:2;

(e) a nucleotide sequence encoding a polypeptide comprising amino acids from about 1 to about 75 in SEQ ID NO:6;

(f) a nucleotide sequence encoding a polypeptide comprising amino acids from about 76 to about 252 in SEQ ID NO:6;

(f) a nucleotide sequence encoding a polypeptide comprising amino acids from about 253 to about 348 in SEQ ID NO:6; and

(g) a nucleotide sequence complementary to any of the nucleotide sequences in (a), (b), (c), (d), (e), or (f).

5           9. An isolated I-FLICE-1 or I-FLICE-2 polypeptide having an amino acid sequence at least 95% identical to a sequence selected from the group consisting of:

(a) amino acids from about 1 to about 480 in SEQ ID NO:2;

(b) amino acids from about 2 to about 480 in SEQ ID NO:2;

10           (c) amino acids from about 1 to about 348 in SEQ ID NO:6;

(d) amino acids from about 2 to about 348 in SEQ ID NO:6;

(e) the amino acid sequence of the I-FLICE-1 polypeptide having the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. 209038;

15           (f) the amino acid sequence of the I-FLICE-2 polypeptide having the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. 209041; and

(g) the amino acid sequence of an epitope-bearing portion of any one of the polypeptides of (a), (b), (c), (d), (e), or (f).

20           10. An isolated antibody that binds specifically to an I-FLICE-1 or I-FLICE-2 polypeptide of claim 9.

11. A method for treating diseases and disorders associated with apoptosis comprising administering to said individual a composition comprising an isolated polypeptide of claim 9.

25           12. A method useful during the diagnosis of diseases and disorders associated with aberrant cell survival in an individual comprising:

(a) measuring I-FLICE-1 or I-FLICE-2 gene expression level in cells or body fluid of said individual;

(b) comparing the I-FLICE-1 or I-FLICE-2 gene expression level of said individual with a standard I-FLICE-1 or I-FLICE-2 gene expression level, whereby an increase or decrease in the I-FLICE-1 or I-FLICE-2 gene expression level of said individual compared to said standard expression level is indicative of disease or disorder associated with aberrant cell survival.

13. An isolated nucleic acid molecule comprising a polynucleotide encoding an I-FLICE-1 or I-FLICE-2 polypeptide wherein, except for at least one conservative amino acid substitution, said polypeptide has a sequence selected from the group consisting of:

(a) a nucleotide sequence encoding a polypeptide comprising amino acids from about 1 to about 480 in SEQ ID NO:2;

(b) a nucleotide sequence encoding a polypeptide comprising amino acids from about 2 to about 480 in SEQ ID NO:2;

(c) a nucleotide sequence encoding a polypeptide comprising amino acids from about 1 to about 348 in SEQ ID NO:6;

(d) a nucleotide sequence encoding a polypeptide comprising amino acids from about 2 to about 348 in SEQ ID NO:6;

(e) a nucleotide sequence encoding a polypeptide having the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. 209038;

(f) a nucleotide sequence encoding a polypeptide having the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. 209041; and

(g) a nucleotide sequence complementary to any of the nucleotide sequences in (a), (b), (c), (d), (e), or (f).

14. An isolated I-FLICE-1 or I-FLICE-2 polypeptide wherein, except for at least one conservative amino acid substitution, said polypeptide has a sequence selected from the group consisting of:

5 (a) amino acids from about 1 to about 480 in SEQ ID NO:2;  
(b) amino acids from about 2 to about 480 in SEQ ID NO:2;  
(c) amino acids from about 1 to about 348 in SEQ ID NO:6;  
(d) amino acids from about 2 to about 348 in SEQ ID NO:6;  
(e) the amino acid sequence of the I-FLICE-1 polypeptide  
10 having the amino acid sequence encoded by the cDNA clone contained in ATCC  
Deposit No. 209038;

(f) the amino acid sequence of the I-FLICE-2 polypeptide  
having the amino acid sequence encoded by the cDNA clone contained in ATCC  
Deposit No. 209041; and

15 (g) the amino acid sequence of an epitope-bearing portion of  
any one of the polypeptides of (a), (b), (c), (d), (e), or (f).

15. An isolated I-FLICE-1 or I-FLICE-2 polypeptide having an amino acid sequence at least 95% identical to a sequence selected from the group consisting of:

20 (a) amino acids from about 1 to about 75 in SEQ ID NO:2;  
(b) amino acids from about 91 to about 171 in SEQ ID NO:2;  
(c) amino acids from about 172 to about 375 in SEQ ID NO:2;  
(d) amino acids from about 376 to about 480 in SEQ ID NO:2;  
(e) amino acids from about 1 to about 75 in SEQ ID NO:6;  
(f) amino acids from about 76 to about 252 in SEQ ID NO:6;  
25 (g) amino acids from about 253 to about 348 in SEQ ID NO:6;

and

(h) the amino acid sequence of an epitope-bearing portion of  
any one of the polypeptides of (a), (b), (c), (d), (e), (f), or (g).

1/11

CGATCGCCCAGCACCAAGTCCGCTTCCAGGCTTTCGGTTTCTTTGCCTCCATCTTGGGTG  
CGCCTTCCCGGCGTCTAGGGGAGCGAAGGCTGAGGTGGCAGCGGCAGGAGAGTCCGGCCG  
CGACAGGACGAACTCCCCACTGGAAAGGATTCTGAAAGAAATGAAGTCAGCCCTCAGAA  
ATGAAGTTGACTGCCTGCTGGCTTTCTGTTGACTGGCCCGGAGCTGTACTGCAAGACCCT  
TGTGAGCTTCCCTAGTCTAAGAGTAGGATGTCTGCTGAAGTCATCCATCAGGTTGAAGAA  
M S A E V I H Q V E E  
GCACTTGATACAGATGAGAAGGAGATGCTGCTCTTTTGTGCCGGGATGTTGCTATAGAT  
A L D T D E K E M L L F L C R D V A I D  
GTGGTTCCACCTAATGTCAGGGACCTTCTGGATATTTTACGGGAAAGAGGTAAGCTGTCT  
V V P P N V R D L L D I L R E R G K L S  
GTCGGGGACTTGGCTGAACTGCTCTACAGAGTGAGGCGATTTGACCTGCTCAAACGTATC  
V G D L A E L L Y R V R R F D L L K R I  
TTGAAGATGGACAGAAAAGCTGTGGAGACCCACCTGCTCAGGAACCTCACCTGTTTCG  
L K M D R K A V E T H L L R N P H L V S  
GACTATAGAGTGCTGATGGCAGAGATTGGTGAGGATTTGGATAAATCTGATGTGTCCTCA  
D Y R V L M A E I G E D L D K S D V S S  
TTAATTTTCTCATGAAGGATTACATGGGCCGAGGCAAGATAAGCAAGGAGAAGAGTTTC  
L I F L M K D Y M G R G K I S K E K S F  
TTGGACCTTGTGGTTGAGTTGGAGAACTAAATCTGGTTGCCCCAGATCAACTGGATTTA  
L D L V V E L E K L N L V A P D Q L D L  
TTAGAAAAATGCCTAAAGAACATCCACAGAATAGACCTGAAGACAAAAATCCAGAAGTAC  
L E K C L K N I H R I D L K T K I Q K Y  
AAGCAGTCTGTTCAAGGAGCAGGGACAAGTTACAGGAATGTTCTCCAAGCAGCAATCCAA  
K Q S V Q G A G T S Y R N V L Q A A I Q  
AAGAGTCTCAAGGATCCTTCAAATAACTTCAGGCTCCATAATGGGAGAAGTAAAGAACAA  
K S L K D P S N N F R L H N G R S K E Q  
AGACTTAAGGAACAGCTTGGCGCTCAACAAGAACCAGTGAAGAAATCCATTGAGGAATCA  
R L K E Q L G A Q Q E P V K K S I Q E S  
GAAGCTTTTTCCTCAGAGCATACCTGAAGAGAGATACAAGATGAAGCAAGCCCTA  
E A F L P Q S I P E E R Y K M K S K P L  
1030 1050 1070

FIG. 1A

2/11

GGAATCTGCCTGATAATCGATTGCATTGGCAATGAGACAGAGCTTCTTCGAGACACCTTC  
G I C L I I D C I G N E T E L L R D T F

1090 1110 1130  
ACTTCCCTGGGCTATGAAGTCCAGAAATCTTGATCTCAGTATGCATGGTATATCCCAG  
T S L G Y E V Q K F L H L S M H G I S Q

1150 1170 1190  
ATTCTTGGCCAATTTGCCTGTATGCCCCAGCACCAGACTACGACAGCTTTGTGTGTGTC  
I L G Q F A C M P E H R D Y D S F V C V

1210 1230 1250  
CTGGTGAGCCGAGGAGGCTCCCAGAGTGTGTATGGTGTGGATCAGACTCACTCAGGGCTC  
L V S R G G S Q S V Y G V D Q T H S G L

1270 1290 1310  
CCCCTGCATCACATCAGGAGGATGTTTCATGGGAGATTTCATGCCCTTATCTAGCAGGGAAG  
P L H H I R R M F M G D S C P Y L A G K

1330 1350 1370  
CCAAAGATGTTTTTATTCAGAACTATGTGGTGTGAGAGGGCCAGCTGGAGGACAGCAGC  
P K M F F I Q N Y V V S E G Q L E D S S

1390 1410 1430  
CTCTTGAGGTGGATGGGCCAGCGATGAAGAATGTGGAATTCAAGGCTCAGAAGCGAGGG  
L L E V D G P A M K N V E F K A Q K R G

1450 1470 1490  
CTGTGCACAGTTCACCGAGAAGCTGACTTCTTCTGGAGCCTGTGTACTGCGGACATGTCC  
L C T V H R E A D F F W S L C T A D M S

1510 1530 1550  
CTGCTGGAGCAGTCTCACAGCTCACCGTCCCTGTACCTGCAGTGCCTCTCCAGAAACTG  
L L E Q S H S S P S L Y L Q C L S Q K L

1570 1590 1610  
AGACAAGAAAGAAAACGCCCACTCCTGGATCTTCACATTGAAGTCAATGGCTACATGTAT  
R Q E R K R P L L D L H I E L N G Y M Y

1630 1650 1670  
GATTGGAACAGCAGAGTTTCTGCCAAGGAGAAATATTATGTCTGGCTGCAGCACACTCTG  
D W N S R V S A K E K Y Y V W L Q H T L

1690 1710 1730  
AGAAAGAACTTATCCTCTCCTACACATAAGAAACCAAAGGCTGGGCGTAGTGGCTCGC  
R K K L I L S Y T \*

1750 1770 1790  
ACCTGTAATCCCAGCACTTTGGGAGGCCAAGGAGGGCGGATCACTTCAGGTCAGGAGTTC  
GAGACCAGCCTGGCCAACATGGTAAACGCTGTCCCTAGTAAGAGTGCAAAAATTAGCTGG

1810 1830 1850  
GTGTGGGTGTGGGTACCTGTGTTCCAGTTACTTGGGAGGCTGAGGTGGGAGGATCTTTT

1870 1890 1910  
GAACCCAGGAGTTCAGGGTCATAGCATGCTGTGATTGTGCCTACGAATAGCCACTGCATA

1930 1950 1970  
CCAACCTGGGCAATATAGCAAGATCCCATCTTTTAAAAAAAAAAAAAAAAAAAAA

1990 2010 2030

FIG. 1B

SUBSTITUTE SHEET (RULE 26)

3/11

1	MD-----F SRNLYDTGEQLDSEDLASLK	huFLICE (U58143)
1	MKSQGGHWYSSSDKNCKVSFREKLLIIDSNLGVQDVENLK	HuMch4 (U60519)
1	MSAEVIH-----QVEEALDTDEKEMUL	HSALZ11Xprotein
1	M-----	HCEBJ50XXprotein
24	FLSLDYIPQRKQEP IKDALM L FQRLQEKRMLEESNLSFLK	huFLICE (U58143)
41	FLCIGLVPNKKLEKSSSASDVFEHLAEDLLSEEDPFFLA	HuMch4 (U60519)
23	FLCRDVAIDVPPNVRD---LLDILRERGLSVGD---LA	HSALZ11Xprotein
2	-----	HCEBJ50XXprotein
64	ELLFRINRLDLLITYLNTRKEEME-RELQTPGRAQISAYR	huFLICE (U58143)
81	ELLYIIRQKKLL-QHLNCTKEEVE-RLL--PTRQRVSLFR	HuMch4 (U60519)
57	ELLYRVRRFDLLKRILKMDRKAVETHLLRNP--HLVSDYR	HSALZ11Xprotein
2	-----	HCEBJ50XXprotein
103	VMLYQIISSEVSRSELRSFKFLQEEISKQKLD DDMNLLDI	huFLICE (U58143)
117	NLLYELSEGIDSENKMDIFLLKDSLPK----TMTSLSF	HuMch4 (U60519)
95	VLMAEIGEDLDKSDVSSLIFLMKDYMGRGKISKEKSFLDL	HSALZ11Xprotein
2	---AEIGEDLDKSDVSSLIFLMKDYMGRGKISKEKSFLDL	HCEBJ50XXprotein
143	F IEMEKRVLIGEGLDILKRVCAGINKSLLKI-INDM---	huFLICE (U58143)
153	LAFLEKQGGKIDEDNLTQLEDLCKTVVPKLLRN-IEKYK--	HuMch4 (U60519)
135	VVELEKLNLVAPDQLDLLEKCLKNIHRIDLKTKIQKYKQS	HSALZ11Xprotein
39	VVELEKLNLVAPDQLDLLEKCLKNIHRIDLKTKIQKYKQS	HCEBJ50XXprotein

FIG.2A

4/11

179	-----EEFSKERSS	LEEGSPDEFS	-----NGEELC	huFLICE (U58143)
190	-----REKAIQIVTPPV	DKEAESYQ	-----GEEEL	HuMch4 (U60519)
175	VQGAGTSYRNVLQAAIQKSLKDPSNNFR	LHNGRSKEQRLK		HSALZ11Xprotein
79	VQGAGTSYRNVLQAAIQKSLKDPSNNFR	-----		HCEBJ50XXprotein

204	GVMTISDS	PREQDS	ISQT	-----LDKVVYQMKSKPRGYC	huFLICE (U58143)
215	----VSQTDVKTFLEALP	-----RAAVYRMNRNHRGLC		HuMch4 (U60519)	
215	EQLGAQQEPVKKSIQ	SEAFLPQSIPEERYKMKSKPLGIC		HSALZ11Xprotein	
107	-----EEPVKKSIQ	SEAFLPQSIPEERYKMKSKPLGIC		HCEBJ50XXprotein	

237	LIINNNHFAKAREKVPKLHS	IRDRNGTHLDAGALITTFEE	huFLICE (U58143)
244	VIIVNNHSFT	-----SLKDRQGTHKDAEILSHVFQW	HuMch4 (U60519)
255	LIIDCIG	-----NETELLRDTFTS	HSALZ11Xprotein
141	LIIDCIG	-----NETELLRDTFTS	HCEBJ50XXprotein

277	LHFEIKPHDDCTVEQIYEILKIYQ-LMDHSNMDCFIQCIL	huFLICE (U58143)
274	LGFTVHIHNNVTKVEMEMVLQKQKCNPAHADGDCFVFCIL	HuMch4 (U60519)
274	LGYE VQKFLHLSMHGISQILGQFACMPEHRDYDSFVCVLV	HSALZ11Xprotein
160	LGYE VQKFLHLSMHGISQILGQFACMPEHRDYDSFVCVLV	HCEBJ50XXprotein

316	SHGDKGI IYGT	DGQEP--PIYELTSQFTGLKCPSLAGKPK	huFLICE (U58143)
314	THGRFGAVYSSDEALI--PIREIMSHFTAQCPRLAGKPK	HuMch4 (U60519)	
314	SRGGSQSVYGV	DQTHSGLPLHHIRRMFMGDSCPYLAGKPK	HSALZ11Xprotein
200	SRGGSQSVYGV	DQTHSGLPLHHIRRMFMGDSCPYLAGKPK	HCEBJ50XXprotein

FIG.2B

SUBSTITUTE SHEET (RULE 26)



5/11

354	VFFIQAC--QG	DNYQKGIPVETD	SEEQPYLEMDLSSPQTR	huFLICE(U58143)
352	LFFIQAC--QGEEIQPS	VSIEADALNPEQAPTS	LDSD---	HuMch4(U60519)
354	MFFIQNYVVSE	GQLEDSSLLEVDGPAMKNVEFKAQKRGLC		HSALZ11Xprotein
240	MFFIQNYVVSD	GQLEDSSLLEVDGPAMKNVEFKAQKRGLC		HCEBJ50XXprotein

392	YIPDEADFL	GMATVNNCM	SYRNPAEGTWY	IQSLCQSLRE	huFLICE(U58143)
387	-IPAEADFL	GLATVPGYMS	FRHVEEGSWY	IQSLCNHKK	HuMch4(U60519)
394	TVHREADFFWSL	CTADMSLLEQSHSSPS	LYLQCL	SQKLRLQ	HSALZ11Xprotein
280	TVHREADFFWSL	CTADMSLLEQSHSSPS	LYLQCL	SQKLRLQ	HCEBJ50XXprotein

432	RCPRGDDILT	ILTEVN	YEVS	NKDDKKNMGKQMPQPTFTL	huFLICE(U58143)
426	LVPRHEDILS	ILTAVN	DDVSRRVDKQGT	KQMPQPAFTL	HuMch4(U60519)
434	--ERKRPL	LDHIEL	NGMYDWN	SRVSAKEKYYVWLQHTL	HSALZ11Xprotein
320	--ERG	-----	TIPG	-----SGITESKDMHFSSLCIL	HCEBJ50XXprotein

471	RKKLVFP	-----	SD	huFLICE(U58143)	
465	RKKLVFPVPL	DALSI		HuMch4(U60519)	
472	RKKLIL	-----	SYTD	HSALZ11Xprotein	
345	-----	L	-----	DVLD	HCEBJ50XXprotein

FIG.2C

6/11




FIG.3

7/11

10	30	50
GCGAGCTTGCAGCCTCACCGACGAGTCTCAACTAAAAGGGACTCCCGGAGCTAGGGGTGG		
70	90	110
GGACTCGGCCTCACACAGTGATTGCCGGCTATTGGACTTTTGTCCAGTGACAGCTGAGAC		
130	150	170
AACAAGGACCACGGGAGGAGGTGTAGGAGAGAAGCGCCGGAACAGGCATCGCCCAGCAC		
190	210	230
CAAGTCCGCTTCCAGGCTTTCGGTTTCTTTGCCTCCATCTTGGGTGCGCCTTCCCGGCGT		
250	270	290
CTAGGGGAGCGAAGGCTGAGGTGGCAGCGGCAGGAGAGTCCGGCCGCGACAGGACGAGTG		
310	330	350
CTGATGGCAGAGATTGGTGAGGATTTGGATAAATCTGATGTGTCCTCATTATTTTCCTC		
M A E I G E D L D K S D V S S L I F L		
370	390	410
ATGAAGGATTACATGGGCCGAGGCAAGATAAGCAAGGAGAAGAGTTTCTTGACCTTG TG		
M K D Y M G R G K I S K E K S F L D L V		
430	450	470
GTTGAGTTGGAGAACTAAATCTGGTTGCCCCAGATCAACTGGATTTATTAGAAAAATGC		
V E L E K L N L V A P D Q L D L L E K C		
490	510	530

FIG.4A

SUBSTITUTE SHEET (RULE 26)

8/11

CTAAAGAACATCCACAGAATAGACCTGAAGACAAAAATCCAGAAGTACAAGCAGTCTGTT  
 L K N I H R I D L K T K I Q K Y K Q S V  
 550 570 590  
 CAAGGAGCAGGGACAAGTTACAGGAATGTTCTCCAAGCAGCAATCCAAAAGAGTCTCAAG  
 Q G A G T S Y R N V L Q A A I Q K S L K  
 610 630 650  
 GATCCTTCAAATAACTTCAGGGAAGAACCAGTGAAGAAATCCATTGAGGAATCAGAAGCT  
 D P S N N F R E E P V K K S I Q E S E A  
 670 690 710  
 TTTTTCCTCAGAGCATACCTGAAGAGAGATACAAGATGAAGAGCAAGCCCCTAGGAATC  
 F L P Q S I P E E R Y K M K S K P L G I  
 730 750 770  
 TGCCTGATAATCGATTGCATTGGCAATGAGACAGAGCTTCTTCGAGACACCTTCACTTCC  
 C L I I D C I G N E T E L L R D T F T S  
 790 810 830  
 CTGGGCTATGAAGTCCAGAAATCTTGCATCTCAGTATGCATGGTATATCCCAGATTCTT  
 L G Y E V Q K F L H L S M H G I S Q I L  
 850 870 890  
 GGCCAATTTGCCTGTATGCCCAGCACCAGACTACGACAGCTTTGTGTGTGTCTGGTG  
 G Q F A C M P E H R D Y D S F V C V L V  
 910 930 950  
 AGCCGAGGAGGCTCCCAGAGTGTGTATGGTGTGGATCAGACTCACTCAGGGCTCCCCCTG  
 S R G G S Q S V Y G V D Q T H S G L P L  
 970 990 1010  
 CATCACATCAGGAGGATGTTTCATGGGAGATTTCATGCCCTTATCTAGCAGGGAAGCCAAAG  
 H H I R R M F M G D S C P Y L A G K P K  
 1030 1050 1070  
 ATGTTTTTTTATTCAGAACTATGTGGTGTGACACGGCCAGCTGGAGGACAGCAGCCTCTTG  
 M F F I Q N Y V V S D G Q L E D S S L L  
 1090 1110 1130  
 GAGGTGGATGGGCCAGCGATGAAGAATGTGGAATTCAAGGCTCAGAAGCGAGGGCTGTGC  
 E V D G P A M K N V E F K A Q K R G L C  
 1150 1170 1190  
 ACAGTTCACCGAGAAGCTGACTTCTTCTGGAGCCTGTGTACTGCGGACATGTCCCTGCTG  
 T V H R E A D F F W S L C T A D M S L L  
 1210 1230 1250  
 GAGCAGTCTCACAGCTCACCGTCCCTGTACCTGCAGTGCCTCTCCAGAAACTGAGACAA  
 E Q S H S S P S L Y L Q C L S Q K L R Q  
 1270 1290 1310  
 GAAAGGGGGACAATTCCCGGAAGTGAATTACAGAGTCAAAGGACATGCATTTTTCAAGC  
 E R G T I P G S G I T E S K D M H F S S  
 1330 1350 1370  
 CTCGGATGCATCTTACTAGATGTCCTATAGGATGGTCATATCAGCTTTATAGGAGAGTAG  
 L G C I L L D V L \*  
 1390 1410 1430  
 CTGTGTCCCTGAATTCTCCCTGACACTGCATGCTCTTATATTTCTCAAGTTTTGACAAT  
 1450 1470 1490  
 TTGATAGGTGAAAAGTGGTATCTGACTGTTTCAGATCTGGAAGGCTTTGTTATATAAACAT  
 1510 1530 1550  
 TTTTTTAATGTTTATTGGCAAGAATACTTTTCTAAGAGAAACATCAGTGAGCTGGTTTTCC  
 1570 1590 1610  
 ATTTAAGCTGAATGAAGCCACAATGTACCTCAAGTATAAGATTAAGTGGCCTTTTTTCAGT  
 1630 1650 1670  
 TGCACCTAATTACAATTTAGAATGATGTTTCTGAGCCACCTGTCAAATGCATTCTGGGC

FIG.4B

9/11

1690 1710 1730  
TGTACCTCTGCGTACCCCAGGAATAAATCTCATGGCCTTCTTTACCTGGCCTCCTTAGTG

1750 1770 1790  
GTGGCCCAGCAGGAAGCGGGGGTTAGAGCAGGAGCCACTCAGCCTTCCAAGATAGATACT

1810 1830 1850  
CCATGGGCGGTTGGTATTACTGGCCTTTTGAGCCCATCCCCATTTGCATAGATGATCCAC

1870 1890 1910  
GTGGGTTATCATCTGGCTGGTATGTTCCCAGAGTGAACTCAGCAGCCCCTTGAGGGAGG

1930 1950 1970  
GGATGGTGGCCATCAGGCCAGAGTATTGCAAGTTAGTTTGGATCATTGCTAAGCAGCTT

1990 2010 2030  
GTGGTGCCCTTCAGAAAGGAACAGTTTCAAAGAACTTTCACATCTGTTGGCTCATTTGCCC

2050 2070 2090  
CTAATGACAGTCTTCTCTTTGATATTTGCATGGCATTAAATTTTGCCTTTCTTGTTTTCT

2110 2130 2150  
CCAGAAAACGCCCACTCCTGGATCTTCACATTGAACTCAATGGCTACATGTATGATTGGA

2170 2190 2210  
ACAGCAGAGTTTCTGCCAAGGAGAAATATTATGTCTGGCTGCAGCACACTCTGAGAAAGA

2230 2250 2270  
AACTTATCTCTCCTACACATAAGAAACCAAAGGCTGGGCGTAGTGGCTCGCACCTGTGA

2290 2310 2330  
TCCCAGCACTTTGGGAGGCCGAGGAGGGCGGATCACTTCAGGTCGGGAGTTCGAGACCAG

2350 2370 2390  
CCTGGCCAGCATGTGAACGCTGTCCCTAGTAGAAGTGCAAAAATTGGCTGGTGTGGGTGT

2410 2430 2450  
GGGTACCCTGTATTCCAGTTGCTTGGGGGGCTGAGGTGGGAGGATCTTTTGACCCAGG

2470 2490 2510  
AGTTCAGGGTCATAGCATGCTGTGATTGTGCCTACGAATAGCCACTGCATACCAACCTGG

2530 2550 2570  
GCAATATAGCAAGATCCCATCTCTTTAAAAAAAAAAAAAAAAAGGACAGGAACCTATCTTAA

2590  
AAAAAAAAAAAAAAAA

FIG.4C

SUBSTITUTE SHEET (RULE 26)

10/11




FIG.5




FIG.6A




FIG.6B

Applicant's or agent's file  
reference number 1488.097PC02

International application No.  
PCT/US98/00969

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM  
(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page 3, line 28.

B. IDENTIFICATION OF DEPOSIT

Further deposits are identified on an additional sheet ☒

Name of depositary institution  
American Type Culture Collection

Address of depositary institution (including postal code and country)

12301 Parklawn Drive  
Rockville, Maryland 20852  
United States of America

Date of deposit  
May 15, 1997

Accession Number  
ATCC 209038

C. ADDITIONAL INDICATIONS (leave blank if not applicable)

This information is continued on an additional sheet ☐

DNA Plasmid 1631454

The applicants hereby request that, until either a Canadian patent has been issued on the basis of the application or the application has been refused, or is abandoned and is no longer subject to reinstatement, or is withdrawn, the furnishing of a sample of deposited biological material referred to in the application only be effected to an independent expert nominated by the Commissioner of Patents.

D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)

E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)

The indications listed below will be submitted to the international Bureau later (specify the general nature of the indications, e.g., "Accession Number of Deposit")

For receiving Office use only

☐ This sheet was received with the international application

Authorized officer

For International Bureau use only

☒ This sheet was received by the International Bureau on:

20 MAY 1998

Authorized officer

Jm



Applicant's or agent's file  
reference number 1488.097PC02

International application No.  
PCT/US98/00969

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM  
(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page 4, line 5.

B. IDENTIFICATION OF DEPOSIT

Further deposits are identified on an additional sheet ☐

Name of depositary institution  
American Type Culture Collection

Address of depositary institution (including postal code and country)

12301 Parklawn Drive  
Rockville, Maryland 20852  
United States of America

Date of deposit  
May 15, 1997

Accession Number  
ATCC 209041

C. ADDITIONAL INDICATIONS (leave blank if not applicable)

This information is continued on an additional sheet ☐

DNA Plasmid 1789630

The applicants hereby request that, until either a Canadian patent has been issued on the basis of the application or the application has been refused, or is abandoned and is no longer subject to reinstatement, or is withdrawn, the furnishing of a sample of deposited biological material referred to in the application only be effected to an independent expert nominated by the Commissioner of Patents.

D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)

E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)

The indications listed below will be submitted to the international Bureau later (specify the general nature of the indications, e.g., "Accession Number of Deposit")

For receiving Office use only

☐ This sheet was received with the international application

Authorized officer

For International Bureau use only

☒ This sheet was received by the International Bureau on:  
20 MAY 1998

Authorized officer

Jm

Applicant's or agent's file  
reference number 1488.097PC02

International application No.

PCT/US 98/00969  
CBA

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM  
(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page 3, line 28.

B. IDENTIFICATION OF DEPOSIT

Further deposits are identified on an additional sheet ☐

Name of depositary institution  
American Type Culture Collection

Address of depositary institution (including postal code and country)

12301 Parklawn Drive  
Rockville, Maryland 20852  
United States of America

Date of deposit  
May 15, 1997

Accession Number  
ATCC 209038

C. ADDITIONAL INDICATIONS (leave blank if not applicable)

This information is continued on an additional sheet ☐

DNA Plasmid 1631454

In respect of those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which the application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28(4) EPC).

D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)

E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)

The indications listed below will be submitted to the international Bureau later (specify the general nature of the indications, e.g., "Accession Number of Deposit")

For receiving Office use only

☒ This sheet was received with the international application

Authorized officer

For International Bureau use only

☐ This sheet was received by the International Bureau on:

Authorized officer

*(DNA Plasmid 1631454)*

#### **FINLAND**

The applicant hereby requests that, until the application has been laid open to public inspection (by the National Board of Patents and Registration), or has been finally decided upon by the National Board of Patents and Registration without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art.

#### **DENMARK**

The applicant hereby request that, until the application has been laid open to public inspection (by the Danish Patent Office), or has been finally decided upon by the Danish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Danish Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Danish Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Danish Patent office or any person approved by the applicant in the individual case.

#### **SWEDEN**

The applicant hereby request that, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Swedish Patent office or any person approved by the applicant in the individual case.

#### **UNITED KINGDOM**

The applicant hereby request that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for international publication of the application.

***(DNA Plasmid 1631454)***


**NETHERLANDS**

The applicant hereby request that until the date of a grant of a Netherlands patent or until the date on which the application is refused or withdrawn or lapsed, the microorganism shall be made available as provided in Rule 31F(1) of the Patent Rules only by the issue of a sample to an expert. The request to this effect must be furnished by the applicant with the Netherlands Industrial Property Office before the date on which the application is made available to the public under Section 22C or Section 25 of the Patents Act of the Kingdom of the Netherlands, whichever of the two dates occurs earlier.

Applicant's or agent's file reference number 1488.097PC02	International application No. <b>PCT/US 98/00969</b> TBA
--	---

**INDICATIONS RELATING TO A DEPOSITED MICROORGANISM**  
(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>4</u> , line <u>5</u> .	
<b>B. IDENTIFICATION OF DEPOSIT</b> <span style="float: right;">Further deposits are identified on an additional sheet <input type="checkbox"/></span>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country)  12301 Parklawn Drive Rockville, Maryland 20852 United States of America	
Date of deposit May 15, 1997	Accession Number ATCC 209041
<b>C. ADDITIONAL INDICATIONS</b> (leave blank if not applicable) <span style="float: right;">This information is continued on an additional sheet <input type="checkbox"/></span>	
DNA Plasmid 1789630  In respect of those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which the application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28(4) EPC).	
<b>D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE</b> (if the indications are not for all designated States)	
<b>E. SEPARATE FURNISHING OF INDICATIONS</b> (leave blank if not applicable)	
The indications listed below will be submitted to the international Bureau later (specify the general nature of the indications, e.g., "Accession Number of Deposit")	

For receiving Office use only	For International Bureau use only
<input checked="" type="checkbox"/> This sheet was received with the international application  Authorized officer	<input type="checkbox"/> This sheet was received by the International Bureau on:  Authorized officer

*(DNA Plasmid 1789630)*

#### **FINLAND**

The applicant hereby requests that, until the application has been laid open to public inspection (by the National Board of Patents and Registration), or has been finally decided upon by the National Board of Patents and Registration without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art.

#### **DENMARK**

The applicant hereby request that, until the application has been laid open to public inspection (by the Danish Patent Office), or has been finally decided upon by the Danish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Danish Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Danish Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Danish Patent office or any person approved by the applicant in the individual case.

#### **SWEDEN**

The applicant hereby request that, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Swedish Patent office or any person approved by the applicant in the individual case.

#### **UNITED KINGDOM**

The applicant hereby request that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for international publication of the application.

***(DNA Plasmid 1789630)***

**NETHERLANDS**

The applicant hereby request that until the date of a grant of a Netherlands patent or until the date on which the application is refused or withdrawn or lapsed, the microorganism shall be made available as provided in Rule 31F(1) of the Patent Rules only by the issue of a sample to an expert. The request to this effect must be furnished by the applicant with the Netherlands Industrial Property Office before the date on which the application is made available to the public under Section 22C or Section 25 of the Patents Act of the Kingdom of the Netherlands, whichever of the two dates occurs earlier.

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 98/00969

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC 6 C12N15/12 C12N9/64 A61K38/48 G01N33/50		
According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) IPC 6 C12N A61K G01N		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	HILLIER L ET AL.: "AC No. AA151642. Homo sapiens cDNA clone 588320" EMBL SEQUENCE DATABASE, 15 December 1996, HEIDELBERG, GERMANY, XP002063883 see the whole document ---	2,8-10, 12,14,15
X	TAKEDA J: "AC No. C05730. EST (expressed sequence tag). Homo sapiens." EMBL SEQUENCE DATABASE, 13 October 1996, HEIDELBERG, GERMANY, XP002063884 see the whole document --- <div style="text-align: center;">-/-</div>	2,9,10, 12,14,15
<div style="display: flex; justify-content: space-between;"> <span><input checked="" type="checkbox"/> Further documents are listed in the continuation of box C.</span> <span><input type="checkbox"/> Patent family members are listed in annex.</span> </div>		
<div style="display: flex;"> <div style="flex: 1;"> <p>* Special categories of cited documents :</p> <p>*A* document defining the general state of the art which is not considered to be of particular relevance</p> <p>*E* earlier document but published on or after the international filing date</p> <p>*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>*O* document referring to an oral disclosure, use, exhibition or other means</p> <p>*P* document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="flex: 1;"> <p>*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>*Z* document member of the same patent family</p> </div> </div>		
Date of the actual completion of the international search  <div style="text-align: center; font-size: 1.2em;">29 April 1998</div>		Date of mailing of the international search report  <div style="text-align: center; font-size: 1.2em;">0 3. 07. 98</div>
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer  <div style="text-align: center; font-size: 1.2em;">Oderwald, H</div>



# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 98/00969

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	CHINNAIYAN A M AND DIXIT V M: "The cell-death machine" CURRENT BIOLOGY, vol. 6, no. 5, 1 May 1996, page 555-562 XP002063885 see page 558, paragraph 2 - page 559, paragraph 6; figure 3; table 1 ---	
A	MUZIO M ET AL: "FLICE, A NOVEL FADD-HOMOLOGOUS ICE/CED-3-LIKE PROTEASE, IS RECRUITED TO THE CD95 (FAS/APO-1) DEATH-INDUCING SIGNALING COMPLEX" CELL, vol. 85, no. 6, 14 June 1996, pages 817-827, XP002037500 cited in the application see abstract; figures 2,3,7 see page 819 - page 822 ---	
P,X	HU S ET AL: "I-FLICE, A NOVEL INHIBITOR OF TUMOR NECROSIS FACTOR RECEPTOR-1- AND CD-95-INDUCED APOPTOSIS" JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 272, no. 28, 11 July 1997, pages 17255-17257, XP002051465 see the whole document ---	1-10, 12-15
P,X	IRMLER M ET AL.: "Inhibition of death receptor signals by cellular FLIP" NATURE, vol. 388, 10 July 1997, pages 190-195, XP002063886 see the whole document -----	1-10, 12-15

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US 98/00969

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
see FURTHER INFORMATION sheet PCT/ISA/210
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Remark : Although claim 11 is directed to a method of treatment of the human/animal body , the search has been carried out and based on the alleged effects of the compound/composition.